


Rasagiline mesylate, MAO-B inhibitor ab120236

2 图像


概述

产品名称	Rasagiline mesylate, MAO-B抑制剂
描述	MAO-B抑制剂
生物学描述	Selective, irreversible monoamine oxidase B (MAO-B) inhibitor, with anti-Parkinson activity. In cell culture (PC-12 and neuroblastoma SH-SY5Y cells) it exhibits neuroprotective and anti-apoptotic activity against several neurotoxins.
CAS编号	161735-79-1
化学结构	 <p>.CH₃SO₃H</p>


性能

化学名称	(1R)-2,3-Dihydro-N-2-propyn-1-yl-1H-Inden-1-amine methanesulfonate
分子量	267.34
分子式	C ₁₂ H ₁₃ N.CH ₃ SO ₃ H
PubChem识别号	3052775
存放说明	Store at +4°C. Store under desiccating conditions. The product can be stored for up to 12 months.
溶解度概述	Soluble in water to 100 mM
处理	<p>Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.</p> <p>Refer to SDS for further information</p> <p>Need more advice on solubility, usage and handling? Please visit our frequently asked questions (FAQ) page for more details.</p>
SMILES	CS(=O)(=O)O.C#CCN[C@@H]2CCc1cccc12
来源	Synthetic

图片



2D chemical structure image of ab120236, Rasagiline mesylate, MAO-B inhibitor



Functional Studies - Rasagiline mesylate, MAO-B inhibitor (ab120236)

ab131457 staining Bcl2 in SK-N-SH cells treated with rasagiline mesylate (ab120236), by ICC/IF. Increase in Bcl2 expression correlates with increased concentration of rasagiline mesylate, as described in literature.

The cells were incubated at 37°C for 24h in media containing different concentrations of ab120236 (rasagiline mesylate) in DMSO, fixed with 100% methanol for 5 minutes at -20°C and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with **ab131457** (1 µg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody (**ab96899**) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team