# abcam

## Product datasheet

## ADAR1 overexpression 293T lysate (whole cell) ab94050

## 2 图像

## 概述

产品名称 ADAR1 overexpression 293T裂解物(whole cell)

常规说明 ab94050 is a 293T cell transfected lysate in which Human ADAR1 has been transiently over-

expressed using a pCMV-ADAR1 plasmid. The lysate is provided in 1X Sample Buffer.

经测试应用 适用于: WB

#### 性能

Mycoplasma free

Yes Liquid

存放说明

形式

Shipped on dry ice. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

存储溶液

Constituents: 0.01% Bromophenol blue, 2.3% Beta mercaptoethanol, 2% Sodium lauryl sulfate,

0.788% Tris HCI, 10% Glycerol (glycerin, glycerine)

背景

Function: Converts multiple adenosines to inosines and creates I/U mismatched base pairs in double-helical RNA substrates without apparent sequence specificity. Has been found to modify more frequently adenosines in AU-rich regions, probably due to the relative ease of melting A/U base pairs as compared to G/C pairs. Functions to modify viral RNA genomes and may be responsible for hypermutation of certain negative-stranded viruses. Edits the messenger RNAs for glutamate receptor (GLUR) subunits by site-selective adenosine deamination. Produces lowlevel editing at the GLUR-B Q/R site, but edits efficiently at the R/G site and HOTSPOT1. Binds to short interfering RNAs (siRNA) without editing them and suppresses siRNA-mediated RNA interference. Binds to ILF3/NF90 and up-regulates ILF3-mediated gene expression. Tissue specificity: Ubiquitously expressed, highest levels were found in brain and lung. Disease: Defects in ADAR are a cause of dyschromatosis symmetrical hereditaria (DSH) [MIM:127400]; also known as reticulate acropigmentation of Dohi. DSH is a pigmentary genodermatosis of autosomal dominant inheritance characterized by a mixture of hyperpigmented and hypopigmented macules distributed on the dorsal parts of the hands and feet. Similarity: Contains

1 A to I editase domain. Contains 2 DRADA repeats. Contains 3 DRBM (double-stranded RNA-


binding) domains. PTM: Sumoylation reduces RNA-editing activity.

#### 应用

## "应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应 <b>用</b>	Ab评论	说 <b>明</b>
WB		Use at an assay dependent dilution.

## 图片



(whole cell) (ab94050)

ab94050 at 15µg/lane on an SDS-PAGE gel.



Western blot - ADAR1 overexpression 293T lysate (whole cell) (ab94050)

All lanes: Anti-ADAR1 antibody (ab88574) at 1/500 dilution

**Lane 1**: ADAR1 overexpression 293T lysate (whole cell) (ab94050)

Lane 2: 293T non-transfected lysate

Lysates/proteins at 25 µg per lane.

#### Secondary

**All lanes :** Goat Anti-mouse IgG (H and L) HRP conjugated at 1/2500 dilution

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors