

### DR6 overexpression 293T lysate (whole cell) ab94106

#### 2 图像

#### 概述

产品名称	DR6 overexpression 293T裂解物(whole cell)
常规说明	ab94106 is a 293T cell transfected lysate in which Human DR6 has been transiently over-expressed using a pCMV-DR6 plasmid. The lysate is provided in 1X Sample Buffer.
经测试应用	适用于: WB

#### 性能

Mycoplasma free	Yes
形式	Liquid
存放说明	Shipped on dry ice. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	Constituents: 0.01% Bromophenol blue, 2.3% Beta mercaptoethanol, 2% Sodium lauryl sulfate, 0.788% Tris HCl, 10% Glycerol (glycerin, glycerine)
背景	Function: May activate NF-kappa-B and promote apoptosis. May activate JNK and be involved in T-cell differentiation. Required for both normal cell body death and axonal pruning. Trophic-factor deprivation triggers the cleavage of surface APP by beta-secretase to release sAPP-beta which is further cleaved to release an N-terminal fragment of APP (N-APP). N-APP binds TNFRSF21 triggering caspase activation and degeneration of both neuronal cell bodies (via caspase-3) and axons (via caspase-6). Tissue specificity: Highly expressed in heart, brain, placenta, pancreas, lymph node, thymus and prostate. Detected at lower levels in lung, skeletal muscle, kidney, testis, uterus, small intestine, colon, spleen, bone marrow and fetal liver. Very low levels were found in adult liver and peripheral blood leukocytes. Similarity: Contains 1 death domain. Contains 4 TNFR-Cys repeats.


#### 应用

**The Abpromise guarantee**      **Abpromise™** 承诺保证使用ab94106于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB		Use at an assay dependent dilution.

## 图片



SDS-PAGE - DR6 overexpression 293T lysate  
(whole cell) (ab94106)

ab94106 at 15µg/lane on an SDS-PAGE gel.



Western blot - DR6 overexpression 293T lysate  
(whole cell) (ab94106)

**All lanes :** Anti-DR6 antibody (**ab89770**) at 1/500 dilution

**Lane 1 :** DR6 overexpression 293T lysate (whole cell) (ab94106)

**Lane 2 :** 293T non-transfected lysate

Lysates/proteins at 25 µg per lane.

### Secondary

**All lanes :** Goat Anti-mouse IgG (H and L) HRP conjugated at 1/2500 dilution

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors