


Anti-58K Golgi protein antibody [58K-9] - Golgi Marker ab27043

★★★★★ [16 Abreviews](#) [60 References](#) [4 图像](#)

概述

产品名称	Anti-58K Golgi蛋白抗体[58K-9] - Golgi Marker
描述	小鼠单克隆抗体[58K-9] to 58K Golgi蛋白- Golgi Marker
宿主	Mouse
经测试应用	适用于: ICC, WB, Flow Cyt (Intra)
种属反应性	与反应: Rat, Human 预测可用于: Hamster, Cow, Dog, Pig, Monkey, African green monkey 
免疫原	Full length native protein (purified). This information is proprietary to Abcam and/or its suppliers.
阳性对照	Rat Liver and HeLa cells. For indirect immunofluorescence: cultured Chinese hamster ovary (CHO) cells For immunoblotting (colorimetric): whole rat liver extract Antigen M.W.: 58 kDa
常规说明	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. Store In the Dark.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine)
纯度	Protein G purified
克隆	单克隆
克隆编号	58K-9

同种型

IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab27043于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC	★★★★☆ (3)	Use a concentration of 5 - 10 µg/ml. Previous batches have worked at the concentration of 1 µg/ml. Our current batch appears to work between 5 and 10 µg/ml. Please see the data below for more details.
WB	★★★★★ (2)	Use a concentration of 1 - 5 µg/ml. Detects a band of approximately 58 kDa (predicted molecular weight: 58 kDa).
Flow Cyt (Intra)		Use 2µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

靶标


功能	Folate-dependent enzyme, that displays both transferase and deaminase activity. Serves to channel one-carbon units from formiminoglutamate to the folate pool. Binds and promotes bundling of vimentin filaments originating from the Golgi.
通路	Amino-acid degradation; L-histidine degradation into L-glutamate; L-glutamate from N-formimidoyl-L-glutamate (transferase route): step 1/1. One-carbon metabolism; tetrahydrofolate interconversion.
疾病相关	Defects in FTCD are the cause of glutamate formiminotransferase deficiency (FIGLU-URIA) [MIM:229100]; also known as formiminoglutamicaciduria (FIGLU-uria). It is an autosomal recessive disorder. Features of a severe phenotype, include elevated levels of formiminoglutamate (FIGLU) in the urine in response to histidine administration, megaloblastic anemia, and mental retardation. Features of a mild phenotype include high urinary excretion of FIGLU in the absence of histidine administration, mild developmental delay, and no hematological abnormalities.
序列相似性	In the C-terminal section; belongs to the cyclodeaminase/cyclohydrolase family. In the N-terminal section; belongs to the formiminotransferase family.
细胞定位	Cytoplasm > cytoskeleton > centrosome > centriole. Golgi apparatus. More abundantly located around the mother centriole.

图片



Flow Cytometry (Intracellular) - Anti-58K Golgi protein antibody [58K-9] - Golgi Marker (ab27043)


Overlay histogram showing HepG2 cells stained with ab27043 (red line). The cells were PFA-fixed and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab27043, 2µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.



Immunocytochemistry - Anti-58K Golgi protein antibody [58K-9] - Golgi Marker (ab27043)

ab27043 staining 58K Golgi protein - Golgi Marker in HeLa cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab27043 at 5µg/ml and [ab6046](#), Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with [ab150117](#), Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and [ab150080](#), Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).


Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Immunocytochemistry - Anti-58K Golgi protein antibody [58K-9] - Golgi Marker (ab27043)

Image courtesy of Dr. Kirk McManus, Univ. of Manitoba/Cancer Care MICB, Canada

ab27043 (1µg/ml, 5µg/ml and 10µg/ml) staining 58K Golgi protein in SK-N-SH cells (green). Cells were fixed in Methanol, permeabilised using 0.5% Triton X100 in PBS and counterstained with DAPI in order to highlight the nucleus (red).



Western blot - Anti-58K Golgi protein antibody [58K-9] - Golgi Marker (ab27043)

All lanes : Anti-58K Golgi protein antibody [58K-9] - Golgi Marker (ab27043) at 1 µg/ml

Lane 1 : Liver (Rat) Tissue Lysate, blocked with 5% BSA

Lane 2 : Liver (Rat) Tissue Lysate, blocked with 3% Milk

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Mouse IgG H&L (HRP) preadsorbed (**ab97040**) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 58 kDa

Exposure time: 3 minutes

Abcam recommends using milk as the blocking agent. Abcam welcomes customer feedback and would appreciate any comments regarding this product and the data presented above.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors