

Anti-6X His tag® antibody [4D11] ab5000

★★★★★ **4 Abreviews** **30 References** **4 图像**

概述

产品名称	Anti-6X His tag®抗体[4D11]
描述	小鼠单克隆抗体[4D11] to 6X His tag®
宿主	Mouse
经测试应用	适用于: Dot blot, ELISA, WB, IP, ICC/IF, Flow Cyt
种属反应性	与反应: Species independent
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
表位	This antibody was shown to recognize an epitope (HHHHHHGS) encoded by the polylinker of several expression vectors (including pDS:RBSII, pQE-30, and pQE-50).
常规说明	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>HIS-TAG® is a trademark of EMD Biosciences, Inc.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Add glycerol to a final volume of 50% for extra stability and aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 6.97% L-Arginine, PBS</p>
纯度	Protein G purified
克隆	单克隆
克隆编号	4D11

骨髓瘤	NS1
同种型	IgG2a
轻链类型	kappa

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab5000于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
AP		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.
WB	★★★★★ (2)	1/1000.
IP	★★★★★ (1)	Use at an assay dependent concentration. See Abreview.
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. ab170191 - Mouse monoclonal IgG2a, is suitable for use as an isotype control with this antibody.

靶标

相关性	The H-H-H-H-H-H motif is used as a tag on many recombinant proteins to facilitate purification. His-tags can be fused to the amino- or carboxy- termini of proteins in transfected or transformed cells.
细胞定位	Depends upon the localization of the parent protein tagged with hexahistidine.

图片



Western blot - Anti-6X His tag® antibody [4D11] (ab5000)

Anti-6X His tag® antibody [4D11] (ab5000) at 1/1000 dilution + 6X His Tagged Protein at 1 µg


Secondary

Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) ([ab65485](#)) at 1/3000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Exposure time: 30 seconds



Immunocytochemistry/ Immunofluorescence - Anti-6X His tag® antibody [4D11] (ab5000)

This image is courtesy of an Abreview submitted by Dr Vladimir Milenkovic

ab5000 staining Human HEK 293 cells transfected with CACNB3-His by ICC/IF. Cells were PFA fixed and permeabilized in 0.5% Triton X-100 prior to blocking in 5% serum for 20 minutes at 25°C. The primary antibody (1/250 in 1% goat serum, 0.1% TX100, 1X PBS) was incubated with the sample for 16 hours at 4°C. An Alexa Fluor® 546 conjugated goat anti-mouse antibody (1/500) was used as the secondary. DAPI was used for nuclei staining (blue).



Western blot - Anti-6X His tag® antibody [4D11] (ab5000)

This image is courtesy of an Abreview submitted by Dr Vladimir Milenkovic


All lanes : Anti-6X His tag® antibody [4D11] (ab5000) at 1/2000 dilution

Lane 1 : Lysates prepared from HEK-293 cells

Lane 2 : Lysates prepared from HEK-293 cells transfected with Beta3-His (CACNB3)

Secondary

All lanes : HRP-conjugated goat anti-mouse IgG polyclonal at 1/5000 dilution



Immunoprecipitation - Anti-6X His tag® antibody [4D11] (ab5000)

This image is courtesy of an Abreview submitted by Dr Vladimir Milenkovic

ab5000 Immunoprecipitate in human HEK293 whole cell lysate. 25µg of cell lysate incubated with primary antibody (undiluted) and matrix (Protein G) for 16 hours at 4°C. For western blotting an undiluted HRP-Goat anti-rabbit IgG polyclonal was used. Beta3-His fusion protein was immunoprecipitated using anti His Ab, and specific beta3a band (56kDa) was detected using anti beta3 Ab

Lane 1. Lysate of HEK293 cells expressing beta3-His (CACNB3)

Lane 2. IP with anti His Ab

Lane 3. Non bound fraction

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors