abcam

Product datasheet

Anti-6X His tag® antibody ab14923

★★★☆☆ 1 Abreviews 15 References 1 图像

概述

产品名称 Anti-6X His tag®抗体

描述 兔多克隆抗体to 6X His tag®

宿主 Rabbit

特异性 ab14923 reacts to recombinant proteins containing the 6X His tag® fused to either the amino or

carboxy terminus in transfected or transformed cells.

经测试应用 适用于: WB

种属反应性 与反应: Species independent

免疫原 Synthetic peptide corresponding to 6X His tag® conjugated to keyhole limpet haemocyanin.

(Peptide available as ab14941, ab14943)

阳性对照 E. coli Positive Control (Escherichia coli) Whole Cell Lysate - expressing 6X His tag protein,

ab2431.

常规说明 HIS-TAG® is a trademark of EMD Biosciences, Inc.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our

1

scientific support team who will be happy to help.

纯**度** Immunogen affinity purified

克隆 多克隆

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab14923于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度:实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★☆☆ (1)	Use a concentration of 0.5 µg/ml.

靶标


相关性 The H-H-H-H motif is used as a tag on many recombinant proteins to facilitate purification.

His-tags can be fused to the amino- or carboxy- termini of proteins in transfected or transformed

cells.

细胞定位 Depends upon the localization of the parent protein tagged with hexahistidine.

图片



Anti-6X His tag® antibody (ab14923) at 0.6 μ g/ml + E. coli Positive Control (Escherichia coli) Whole Cell Lysate - expressing 6X His tag protein (ab2431) at 10 μ g

Secondary

Alexa Fluor Goat polyclonal to Rabbit IgG at 1/10000 dilution

Performed under reducing conditions.

Observed band size: 17 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors