

Anti-ABCC10 antibody [M7I-3] ab91451

8 References [1 图像](#)

概述

产品名称	Anti-ABCC10抗体[M7I-3]
描述	大鼠单克隆抗体[M7I-3] to ABCC10
宿主	Rat
经测试应用	适用于: ICC
种属反应性	与反应: Human
免疫原	Recombinant fragment within Human ABCC10 aa 150-300 (internal sequence). The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please contact our Scientific Support team to discuss your requirements.



[Run BLAST with](#)



[Run BLAST with](#)

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	pH: 7.3 Preservative: 0.1% Sodium azide Constituent: 0.7% BSA
	Serum free tissue culture supernatant
纯度	Tissue culture supernatant
克隆	单克隆
克隆编号	M7I-3

同种型

IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用 ab91451 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
ICC		1/20 - 1/50. Acetone fixed cytospin preparations.

靶标

功能	ATP-dependent transporter probably involved in cellular detoxification through lipophilic anion extrusion.
组织特异性	Isoform 1 is specifically expressed in spleen. Isoform 2 is more widely expressed.
序列相似性	Belongs to the ABC transporter superfamily. ABCC family. Conjugate transporter (TC 3.A.1.208) subfamily. Contains 2 ABC transmembrane type-1 domains. Contains 2 ABC transporter domains.
细胞定位	Cell membrane.

图片



ab91451 at 1/20 dilution staining ABCC10 in ABCC10 transfected HEK cells by Immunocytochemistry.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors