


Anti-ACK1 antibody ab65108

★ ★ ★ ★ ★ [1 Abreviews](#) [3 References](#) [2 图像](#)

概述

产品名称	Anti-ACK1抗体
描述	兔多克隆抗体to ACK1
宿主	Rabbit
经测试应用	适用于: ICC/IF, WB
种属反应性	与反应: Rat, Human 预测可用于: Mouse, Cow, Dog, Rhesus monkey 
免疫原	Synthetic peptide conjugated to KLH derived from within residues 1000 to the C-terminus of Human ACK1.参阅Abcam的专有抗源政策(Peptide available as ab74487 .)
阳性对照	This antibody gave a positive signal in the following Whole Cell Lysates: HeLa, HepG2, PC12, SHSY-5Y, T24/83, HT 1080
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS
纯度	Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help. Immunogen affinity purified

克隆	多克隆
同种型	IgG

应用


The Abpromise guarantee **Abpromise™**承诺保证使用ab65108于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★☆☆☆☆ (1)	Use a concentration of 1 µg/ml.
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 120 kDa (predicted molecular weight: 114 kDa).

靶标

功能	Non-receptor tyrosine kinase that regulates the activity of a number of proteins by tyrosine phosphorylation especially proteins critical for cell survival, cell growth, and proliferation. Activates AKT1 by phosphorylating it on 'Tyr-176' resulting in its activation. Phosphorylates AR on 'Tyr-267' and 'Tyr-363' and promotes its recruitment to the androgen-responsive enhancers (AREs). Phosphorylates WWOX on 'Tyr-287'. Downstream effector of CDC42 which mediates CDC42-dependent cell migration via phosphorylation of BCAR1. Binds to both poly- and mono-ubiquitin and regulates ligand-induced degradation of EGFR. Participates in clathrin-mediated endocytosis. May be involved both in adult synaptic function and plasticity and in brain development.
组织特异性	The Tyr-284 phosphorylated form shows a significant increase in expression in breast cancers during the progressive stages i.e. normal to hyperplasia (ADH), ductal carcinoma in situ (DCIS), invasive ductal carcinoma (IDC) and lymph node metastatic (LNMM) stages. It also shows a significant increase in expression in prostate cancers during the progressive stages.
序列相似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. Contains 1 CRIB domain. Contains 1 protein kinase domain. Contains 1 SH3 domain. Contains 1 UBA domain.
结构域	The EBD (EGFR-binding domain) domain is necessary for interaction with EGFR. The SAM-like domain is necessary for NEDD4-mediated ubiquitination. Promotes membrane localization and dimerization to allow for autophosphorylation. The UBA domain binds both poly- and mono-ubiquitin.
翻译后修饰	Autophosphorylation regulates kinase activity. Phosphorylation on Tyr-518 is required for interaction with SRC. Ubiquitinated by NEDD4. Its EGF-induced degradation is EGFR activation-dependent and is processed by lysosomes, not proteasomes.
细胞定位	Cell membrane. Nucleus. Endosome. Cell junction > adherens junction. Cytoplasmic vesicle membrane. The Tyr-284 phosphorylated form is expressed both in the membrane and nucleus. Co-localizes with EGFR on the endosomes.



All lanes : Anti-ACK1 antibody (ab65108) at 1 µg/ml

Lane 1 : HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate

Lane 2 : HepG2 (Human hepatocellular liver carcinoma cell line) Whole Cell Lysate

Lane 3 : PC12 (Rat adrenal pheochromocytoma cell line) Whole Cell Lysate

Lane 4 : SHSY-5Y (Human neuroblastoma cell line) Whole Cell Lysate

Lane 5 : T24/83 (Human bladder carcinoma) Whole Cell Lysate

Lane 6 : HT 1080 (Human fibrosarcoma) Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 114 kDa

Observed band size: 120 kDa

Exposure time: 4 minutes



Immunocytochemistry/ Immunofluorescence - Anti-ACK1 antibody (ab65108)

ICC/IF image of ab65108 stained HeLa cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab65108, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors