

Anti-Actin antibody [EPR16769] - BSA and Azide free ab219733

重组 RabMAb

6 References **9 图像**

概述

产品名称	Anti-Actin抗体[EPR16769] - BSA and Azide free
描述	兔单克隆抗体[EPR16769] to Actin - BSA and Azide free
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF, IP, Flow Cyt (Intra)
种属反应性	与反应: Mouse, Rat, Chicken, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa, 293T, C6, RAW 264.7, PC-12, NIH/3T3 and UMNSAH/DF-1 whole cell lysates; Human skeletal muscle, fetal spleen, fetal brain, fetal heart, fetal kidney and cardiac muscle lysates; Mouse and Rat brain, heart, kidney and spleen lysates. IHC-P: Human prostate hyperplasia, Mouse skeletal muscle and Rat skeletal muscle tissues. ICC/IF: NIH/3T3 cells. IP: NIH/3T3 whole cell extract. Flow Cyt (intra): HeLa cells
常规说明	<p>ab219733 is the carrier-free version of ab179467.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR16769
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用ab219733于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use at an assay dependent concentration. Detects a band of approximately 42 kDa (predicted molecular weight: 42 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
Flow Cyt (Intra)		Use at an assay dependent concentration.

靶标

功能	Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.
疾病相关	Defects in ACTA1 are the cause of nemaline myopathy type 3 (NEM3) [MIM:161800]. A form of nemaline myopathy. Nemaline myopathies are muscular disorders characterized by muscle weakness of varying severity and onset, and abnormal thread-or rod-like structures in muscle fibers on histologic examination. The phenotype at histological level is variable. Some patients present areas devoid of oxidative activity containing (cores) within myofibers. Core lesions are unstructured and poorly circumscribed. Defects in ACTA1 are a cause of myopathy congenital with excess of thin myofilaments (MPCETM) [MIM:161800]. A congenital muscular disorder characterized at histological level by

areas of sarcoplasm devoid of normal myofibrils and mitochondria, and replaced with dense masses of thin filaments. Central cores, rods, ragged red fibers, and necrosis are absent. Defects in ACTA1 are a cause of congenital myopathy with fiber-type disproportion (CFTD) [MIM:255310]; also known as congenital fiber-type disproportion myopathy (CFTDM). CFTD is a genetically heterogeneous disorder in which there is relative hypotrophy of type 1 muscle fibers compared to type 2 fibers on skeletal muscle biopsy. However, these findings are not specific and can be found in many different myopathic and neuropathic conditions.


序列相似性

Belongs to the actin family.

细胞定位


Cytoplasm > cytoskeleton.

图片




Immunocytochemistry/ Immunofluorescence analysis of NIH/3T3 (Mouse embryonic fibroblast) cells labeling Actin with Purified **ab179467** at 1:100 dilution (6.98 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 dilution (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).



Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelia cell) cells labeling Actin with Purified **ab179467** at 1:100 dilution (6.98 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 dilution (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).



Flow Cytometry (Intracellular) - Anti-Actin antibody
[EPR16769] - BSA and Azide free (ab219733)

Intracellular Flow Cytometry analysis of HeLa (human cervix adenocarcinoma) cells labelling Actin with purified **ab179467** at 1/70 (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. An Alexa Fluor®488-conjugated goat anti-rabbit IgG (1/2000) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Actin antibody
[EPR16769] - BSA and Azide free (ab219733)

Immunohistochemical analysis of paraffin-embedded Human prostate hyperplasia tissue labeling Actin with **ab179467** at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm staining on smooth muscle cells is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Actin antibody [EPR16769] - BSA and Azide free (ab219733)

Immunohistochemical analysis of paraffin-embedded Mouse skeletal muscle tissue labeling Actin with **ab179467** at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm staining is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Actin antibody [EPR16769] - BSA and Azide free (ab219733)

Immunohistochemical analysis of paraffin-embedded Rat skeletal muscle tissue labeling Actin with **ab179467** at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm staining is observed. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).

Heat mediated antigen retrieval was performed with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-Actin antibody [EPR16769] - BSA and Azide free (ab219733)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryo fibroblast cells) cells labeling Actin with **ab179467** at 1/50 dilution, followed by Goat **anti-rabbit Alexa Fluor® 488** (IgG) (**ab150077**) secondary antibody at 1/200 dilution (green). Cytoplasm staining on NIH/3T3 cell line is observed. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution and **ab150120** (goat **anti-mouse AlexaFluor®594** secondary) at 1/500 dilution (red).

The negative controls are as follows;

1. **ab179467** at 1/50 dilution followed by **ab150120** (goat **anti-mouse AlexaFluor®594** secondary) at 1/500 dilution.
2. **ab7291** (anti-Tubulin mouse mAb) at 1/500 dilution followed by **ab150077** (goat **anti-rabbit Alexa Fluor®488** (IgG H&L) at 1/200 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).



Immunoprecipitation - Anti-Actin antibody [EPR16769] - BSA and Azide free (ab219733)

Actin was immunoprecipitated from 1mg of NIH/3T3 (Mouse embryo fibroblast cells) whole cell extract with **ab179467** at 1/40 dilution. Western blot was performed from the immunoprecipitate using **ab179467** at 1/1000 dilution. Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated was used as secondary antibody at 1/1000 dilution. Lane 1: NIH/3T3 whole cell extract. Lane 2: PBS instead of NIH/3T3 whole cell extract.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab179467**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Actin antibody [EPR16769] - BSA and Azide free (ab219733)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors