

Anti-Acyloxyacyl Hydrolase antibody ab233405

12 图像

概述

产品名称	Anti-Acyloxyacyl Hydrolase 抗体
描述	兔多克隆抗体 to Acyloxyacyl Hydrolase
宿主	Rabbit
经测试应用	适用于: IHC-P, WB
种属反应性	与反应: Human, Pig
免疫原	Recombinant fragment (His-T7-tag) corresponding to Human Acyloxyacyl Hydrolase aa 1-300. (Expressed in E.coli). Database link: P28039

 [Run BLAST with](#)

 [Run BLAST with](#)

阳性对照	IHC-P: Human appendix, breast cancer, prostate cancer, glioma, liver and stomach tissues. WB: Human serum; Pig stomach, small intestine and spleen lysates; Recombinant human Acyloxyacyl Hydrolase protein.
------	--

常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
------	---

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.011% Proclin 300 Constituents: 55.77% Glycerol (glycerin, glycerine), 44.219% PBS
纯度	Immunogen affinity purified
纯化说明	ab233405 was purified by antigen-specific affinity chromatography followed by Protein A affinity chromatography.

克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab233405于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use a concentration of 5 - 20 µg/ml.
WB		Use a concentration of 0.2 - 20 µg/ml. Predicted molecular weight: 65 kDa.

靶标


功能 Removes the secondary (acyloxyacyl-linked) fatty acyl chains from the lipid A region of bacterial lipopolysaccharides.

序列相似性 Contains 1 saposin B-type domain.

翻译后修饰 Both subunits contain a number of cysteine residues that may form disulfide bridges.


细胞定位 Secreted.

图片




Paraffin-embedded human appendix tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)




Paraffin-embedded human breast cancer tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)




Paraffin-embedded human prostate cancer tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)




Paraffin-embedded human glioma tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)




Paraffin-embedded human liver tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Paraffin-embedded human stomach tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 $\mu\text{g}/\text{ml}$ in immunohistochemical analysis. DAB staining.



Western blot - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Anti-Acyloxyacyl Hydrolase antibody (ab233405) at 2 $\mu\text{g}/\text{ml}$ + Recombinant human Acyloxyacyl Hydrolase protein

Predicted band size: 65 kDa



Western blot - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Anti-Acyloxyacyl Hydrolase antibody (ab233405) at 2 μ g/ml + Pig stomach lysate

Predicted band size: 65 kDa



Western blot - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Anti-Acyloxyacyl Hydrolase antibody (ab233405) at 2 μ g/ml + Human serum

Predicted band size: 65 kDa



Western blot - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Anti-Acyloxyacyl Hydrolase antibody (ab233405) at 2 μ g/ml + Pig small intestine lysate


Predicted band size: 65 kDa



Western blot - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Anti-Acyloxyacyl Hydrolase antibody (ab233405) at 2 µg/ml + Pig spleen lysate

Predicted band size: 65 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acyloxyacyl Hydrolase antibody (ab233405)

Formalin-fixed, paraffin-embedded human pancreas tissue stained for Acyloxyacyl Hydrolase using ab233405 at 20 µg/ml in immunohistochemical analysis. DAB staining.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors