abcam

Product datasheet

مل سما

Anti-ADAM17 antibody ab28233

★★★★★ <u>5 Abreviews</u> <u>5 References</u>

199. 202			
产品名称	Anti-ADAM17 抗体		
描述	兔多克隆抗体to ADAM17		
宿主	Rabbit		
经 测 试应 用	适用于: IP, ICC/IF, WB		
种属反应性	与反 应: Rat, Human, Pig		
	预测可用于: Mouse 🔺		
免疫原	Synthetic peptide based on the amino end of the catalytic domain of human ADAM-17. 参阅 Abcam 的 专 有抗源政策 (Peptide available as <u>ab41216</u> .)		
常 规说 明	The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.		
	If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As		
性能			
形式	Liquid		
存 放 说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.		
存储溶液	Preservative: 0.05% Sodium azide Constituent: 50% Glycerol		
纯 度	Immunogen affinity purified		
克隆	多克隆		
同种型	lgG		

应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP	★ ★ ★ ★ ★ <u>(1)</u>	Use at an assay dependent concentration.
ICC/IF	★ ★ ★ ★ ★ (1)	Use at an assay dependent concentration.
WB	★ ★ ★ ☆ ☆ (2)	Use at an assay dependent concentration. Predicted molecular weight: 93 kDa. ab28233 recognizes a 120 kDa band in reduced Western blots of conditioned media or cell lysates, which is converted to a 55- 60 kDa band. EDTA/EGTA treatment of tissues or lysates is required to see latent zymogen. Note: 1/1,000 when using colorimetric substrates such as BCIP/NBT, and 1/5,000 for

靶标

Cleaves the membrane-bound precursor of TNF-alpha to its mature soluble form. Responsible for the proteolytical release of soluble JAM3 from endothelial cells surface. Responsible for the proteolytic release of several other cell-surface proteins, including p75 TNF-receptor, interleukin 1 receptor type II, p55 TNF-receptor, transforming growth factor-alpha, L-selectin, growth hormone receptor, MUC1 and the amyloid precursor protein. Also involved in the activation of Notch pathway.		
Ubiquitously expressed. Expressed at highest levels in adult heart, placenta, skeletal muscle, pancreas, spleen, thymus, prostate, testes, ovary and small intestine, and in fetal brain, lung, liver and kidney.		
Contains 1 disintegrin domain. Contains 1 peptidase M12B domain.		
Must be membrane anchored to cleave the different substrates. The cytoplasmic domain is not required for the this activity. Only the catalytic domain is essential to shed TNF and p75 TNFR. The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.		
The precursor is cleaved by a furin endopeptidase. Phosphorylated. Stimulation by growth factor or phorbol 12-myristate 13-acetate induces phosphorylation of Ser-819 but decreases phosphorylation of Ser-791.		
Membrane.		

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <u>https://www.abcam.cn/abpromise</u> or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors