abcam

Product datasheet

Anti-Adenine Nucleotide Translocase Immunocapture antibody [5F51BB5AG7] ab109864

5 References 2 图像

概述

产品名称 Anti-Adenine Nucleotide Translocase Immunocapture抗体[5F51BB5AG7]

描述 小鼠单克隆抗体[5F51BB5AG7] to Adenine Nucleotide Translocase Immunocapture

宿主 Mouse

 经测试应用
 适用于: IP, IHC-P

 种属反应性
 与反应: Cow. Human

免疫原 Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.

阳性对照 Bovine and Human heart mitochondria This antibody gave a positive result in IHC in the following

FFPE tissue: Human normal heart muscle.

常规说明

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or

conjugation for your experiments, please contact orders@abcam.com.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

Product was previously marketed under the MitoSciences sub-brand.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C. Do Not Freeze.

存储溶液 pH: 7.5

Preservative: 0.02% Sodium azide Constituent: HEPES buffered saline

纯**化说明** ab109864 was produced in vitro using hybridomas grown in serum-free medium, and then

purified by biochemical fractionation.

1

克隆 单克隆

克隆编号 5F51BB5AG7

同种型 IgG1

轻链类型 kappa

应用

The Abpromise guarantee Abpromise™承诺保证使用ab109864于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		Use at 100 µg/mg of lysate.
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标


相关性

ANT is a reversible transporter of ATP and ADP. During oxidative phosphorylation (OXPHOS) the protein exchanges ATP out for ADP in, however, in the absence of OXPHOS ANT works in reverse to maintain mitochondrial membrane potential. A subset of patients with heart related diseases have been shown to harbor mutations in the ANT gene or to develop an autoimmune reaction against the protein. There are a number of proteins found in mitochondria that, under circumstances favoring apoptosis, coalesce to form the so-called permeability transition pore. The proteins identified as a part of the PTP include cytosolic hexokinase, outer membrane porin (also called the voltage dependent anion channel or VDAC), adenylate cyclase in the intermembrane space, the adenine nucleotide transporter (ANT) and the peripheral benzodiazepine receptor protein of the inner membrane along with cyclophilin D of the matrix space. Hexokinase, porin and ANT all occur as isoforms and it remains to be determined whether there is specificity of these different forms for the PTP.

细胞定位

Mitochondrial

图片




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Adenine Nucleotide

Translocase Immunocapture antibody

[5F51BB5AG7] (ab109864)

IHC image of Adenine Nucleotide Translocase staining in Human normal heart muscle formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab109864, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Translocase Immunocapture antibody

[5F51BB5AG7] (ab109864)

Immunocapture of Adenine Nucleotide Translocase from Bovine heart mitochondria using ab109864 at a concentration of 100 $\mu g/mg$ lysate.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors