abcam

Product datasheet

Alexa Fluor® 488 Anti-Caldesmon/CDM antibody [E89] ab208116



重组 RabMAb

2 图像

概述

产品名称 Alexa Fluor® 488荧光Anti-Caldesmon/CDM抗体[E89]

描述 Alexa Fluor® 488荧光兔单克隆抗体[E89] to Caldesmon/CDM

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

经测试应用 适用干: ICC/IF 种属反应性 与反应: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 ICC/IF: HeLa cells.

常规说明 Our RabMAb® technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to **RabMAb® patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

纯**度** Protein A purified

 克隆
 单克隆

 克隆编号
 E89

 同种型
 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab208116于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/100. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

功能 Actin- and myosin-binding protein implicated in the regulation of actomyosin interactions in

smooth muscle and nonmuscle cells (could act as a bridge between myosin and actin filaments). Stimulates actin binding of tropomyosin which increases the stabilization of actin filament structure. In muscle tissues, inhibits the actomyosin ATPase by binding to F-actin. This inhibition is attenuated by calcium-calmodulin and is potentiated by tropomyosin. Interacts with actin, myosin, two molecules of tropomyosin and with calmodulin. Also play an essential role during

cellular mitosis and receptor capping.

组织特异性 High-molecular-weight caldesmon (isoform 1) is predominantly expressed in smooth muscles,

whereas low-molecular-weight caldesmon (isoforms 2, 3, 4 and 5) are widely distributed in non-

muscle tissues and cells. Not expressed in skeletal muscle or heart.

序列相似性 Belongs to the caldesmon family.

结**构域** The N-terminal part seems to be a myosin/calmodulin-binding domain, and the C-terminal a

tropomyosin/actin/calmodulin-binding domain. These two domains are separated by a central

helical region in the smooth-muscle form.

翻译后修饰 In non-muscle cells, phosphorylation by CDK1 during mitosis causes caldesmon to dissociate

from microfilaments. Phosphorylation reduces caldesmon binding to actin, myosin, and


calmodulin as well as its inhibition of actomyosin ATPase activity. Phosphorylation also occurs in both quiescent and dividing smooth muscle cells with similar effects on the interaction with actin

and calmodulin and on microfilaments reorganization.

细胞定位 Cytoplasm > cytoskeleton. Cytoplasm > myofibril. On thin filaments in smooth muscle and on

stress fibers in fibroblasts (nonmuscle).

图片




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Caldesmon/CDM antibody [E89] (ab208116)

ab208116 staining Caldesmon/CDM in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab208116 at 1/100 dilution (shown in green) and <u>ab195889</u>, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 4% formaldehyde (10 min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors