abcam

Product datasheet

Alexa Fluor® 488 Anti-S100A9 antibody [EPR3555] ab200972



重组 RabMAb

2 图像

概述

产品名称 Alexa Fluor® 488荧光Anti-S100A9抗体[EPR3555]

描述 Alexa Fluor® 488荧光兔单克隆抗体[EPR3555] to S100A9

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

适用于: ICC/IF 经测试应用 种属反应性 与反应: Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 ICC/IF: Jurkat cells. Flow Cyt: Jurkat cells.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 常规说明 monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

> Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

性能

形式

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

1

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

纯**度** Protein A purified

 克隆
 单克隆

 克隆编号
 EPR3555

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab200972于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/50. This product gave a positive signal in Jurkat cells fixed with 100% methanol (5 min).

靶标

功能 Calcium-binding protein. Has antimicrobial activity towards bacteria and fungi. Important for

resistance to invasion by pathogenic bacteria. Up-regulates transcription of genes that are under the control of NF-kappa-B. Plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS) (By similarity). Promotes tubulin polymerization when unphosphorylated. Promotes phagocyte migration and infiltration of granulocytes at sites of wounding. Plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Extracellular calprotectin binds to target cells and promotes apoptosis. Antimicrobial and proapoptotic activity is inhibited by zinc

ions.

组织特异性 Expressed by macrophages in acutely inflammed tissues and in chronic inflammation. Detected

in peripheral blood leukocytes, in neutrophils and granulocytes. Detected at sites of vascular inflammation (at protein level). Also expressed in epithelial cells constitutively or induced during

dermatoses.

序列相似性 Belongs to the S-100 family.

Contains 2 EF-hand domains.


翻译后修饰 Phosphorylated. Phosphorylation inhibits activation of tubulin polymerization.

细胞定位 Secreted. Cytoplasm. Cytoplasm > cytoskeleton. Cell membrane. Associates with tubulin

filaments in activated monocytes. Targeted to the cell surface upon calcium influx. Released from blood leukocytes upon exposure to CSF2/GM-CSF, bacterial lipopolysaccharide (LPS) and during inflammatory processes. Serum levels are high in patients suffering from chronic


inflammation.

图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-S100A9 antibody [EPR3555] (ab200972)

ab200972 staining S100A9 in Jurkat cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab200972 at 1/50 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors