abcam

Product datasheet

Alexa Fluor® 488 Anti-TSG101 antibody [EPR7130(B)] ab207663



重组 RabMAb

1 References 2 图像

概述

免疫原

产品名称 Alexa Fluor® 488荧光Anti-TSG101抗体[EPR7130(B)]

描述 Alexa Fluor® 488荧光兔单克隆抗体[EPR7130(B)] to TSG101

宿主 Rabbit

偶联物 Alexa Fluor® 488. Ex: 495nm, Em: 519nm

适用干: ICC/IF 经测试应用 种属反应性 与反应: Human

预测可用于: Mouse, Rat 🔷

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ICC/IF: Jurkat cells. 阳性对照

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

1

outlicensing@thermofisher.com.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR7130(B)

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab207663于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度:实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|--------|------|---|
| ICC/IF | | 1/100. This product gave a positive signal in Jurkat cells fixed with 4% formaldehyde (10 min). |

| 靶 | 7 | ķ | Ŕ |
|---|---|---|---|
| | | | |

功能 Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to

ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55. May be involved in cell growth and differentiation. Acts as a negative growth regulator. Involved in the budding of many viruses through an interaction with viral proteins that contain a late-budding motif P-[ST]-A-

P. This interaction is essential for viral particle budding of numerous retroviruses.

组织特异性 Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas.

序列相似性 Belongs to the ubiquitin-conjugating enzyme family. UEV subfamily.

Contains 1 SB (steadiness box) domain.
Contains 1 UEV (ubiquitin E2 variant) domain.

结**构域** The UEV domain is required for the interaction of the complex with ubiquitin. It also mediates the

interaction with PTAP/PSAP motifs of HIV-1 P6 protein and human spumaretrovirus Gag protein.

The coiled coil domain may interact with stathmin.


The UEV domain binds ubiquitin and P-[ST]-A-P peptide motif independently.

翻译后修饰 Monoubiguitinated at multiple sites by LRSAM1 and by MGRN1. Ubiquitination inactivates it,

possibly by regulating its shuttling between an active membrane-bound protein and an inactive

Cytoplasm. Membrane. Nucleus. Late endosome membrane. Mainly cytoplasmic. Membrane-associated when active and soluble when inactive. Depending on the stage of the cell cycle, detected in the nucleus. Colocalized with CEP55 in the midbody during cytokinesis.

图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-TSG101 antibody [EPR7130(B)] (ab207663)

ab207663 staining TSG101 in Jurkat cells. The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab207663 at 1/100 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors