# abcam

## Product datasheet

# Alexa Fluor® 647 Anti-RENT1/hUPF1 antibody [EPR4681] ab201265



重组 RabMAb

## 2 图像

### 概述

产品名称 Alexa Fluor® 647荧光Anti-RENT1/hUPF1抗体[EPR4681]

描述 Alexa Fluor® 647荧光兔单克隆抗体[EPR4681] to RENT1/hUPF1

宿主 Rabbit

偶联物 Alexa Fluor® 647. Ex: 652nm, Em: 668nm

经测试应用 适用干: ICC/IF 种属反应性 与反应: Human

预测可用于: Mouse 🗥

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 ICC/IF: HeLa cells

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 常规说明

monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor<sup>®</sup> is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific Company. The Alexa Fluor<sup>®</sup> dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

## 性能

形式 Liquid

**存放说明** Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Stable for 12 months at -20°C. Store In the Dark.

**存储溶液** pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), PBS, 1% BSA

纯**度** Protein A purified

**同种型** IgG

#### 应用

# The Abpromise guarantee Abpromise™承诺保证使用ab201265于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/100.

#### 靶标

功能 Plays a role in replication-dependent histone mRNA degradation at the end of phase S. Part of a

post-splicing multiprotein complex. Involved in nonsense-mediated decay (NMD) as part of the SMG1C complex, a mRNA surveillance complex that recognizes and degrades mRNAs containing premature translation termination codons (PTCs). The complex probably acts by associating with ribosomes during tranlation termination on mRNPs. If an exon junction complex (EJC) is located 50-55 or more nucleotides downstream from the termination codon, RENT1 is phosphorylated by SMG1, triggering nonsense-mediated decay (NMD). Essential for embryonic

viability.

组织特异性 Ubiquitous.

序列相似性 Belongs to the DNA2/NAM7 helicase family.

Contains 1 C2H2-type zinc finger.

结构域 The [ST]-Q motif constitutes a recognition sequence for kinases from the Pl3/Pl4-kinase family.


翻译后修饰 Phosphorylated by SMG1; required for formation of mRNA surveillance complexes.

Phosphorylated upon DNA damage, probably by ATM or ATR.

细胞定位 Cytoplasm. Cytoplasm > P-body. Hyperphosphorylated form is targeted to the P-body, while

unphosphorylated protein is distributed throughout the cytoplasm.

### 图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-RENT1/hUPF1 antibody [EPR4681] (ab201265)

ab201265 staining RENT1/hUPF1 in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab201265 at 1/100 dilution (shown in red) and <a href="mailto:ab195887">ab195887</a>, Mouse monoclonal to alpha Tubulin (Alexa Fluor<sup>®</sup> 488) at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 4% formaldehyde (10 min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

• We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors