# abcam

# Product datasheet

# Alexa Fluor® 647 Anti-TAK1 antibody [EPR5984] ab199473





重组 RabMAb

4 图像

#### 概述

产品名称 Alexa Fluor® 647荧光Anti-TAK1抗体[EPR5984]

描述 Alexa Fluor® 647荧光兔单克隆抗体[EPR5984] to TAK1

宿主 Rabbit

偶联物 Alexa Fluor® 647. Ex: 652nm, Em: 668nm

适用于: ICC/IF 经测试应用

种属反应性 与反应: Mouse, Human

预测可用于: Rat 📤

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 ICC/IF: differentiated Neuro2a and MCF7 cells, HAP1 cells (WT and MAP3K7-KO).

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit 常规说明

monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor<sup>®</sup> is a registered trademark of Molecular Probes, Inc., a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

性能

形式

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

**存储溶液** pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), PBS, 1% BSA

纯**度** Protein A purified

同种型 lgG

#### 应用

# The Abpromise guarantee Abpromise™承诺保证使用ab199473于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/100. This product gave a positive signal in differentiated Neuro2a and MCF7 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

ध्या	丰二
牝	仦

功能 Component of a protein kinase signal transduction cascade. Mediator of TRAF6 and TGF-beta

signal transduction. Activates IKBKB and MAPK8 in response to TRAF6 signaling. Stimulates NF-kappa-B activation and the p38 MAPK pathway. In osmotic stress signaling, plays a major

role in the activation of MAPK8/JNK, but not that of NF-kappa-B.

序列相似性 Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase

kinase subfamily.

Contains 1 protein kinase domain.

翻译后修饰 Association with TAB1/MAP3K7IP1 promotes autophosphorylation and subsequent activation.

Association with TAB2/MAP3K7IP2, itself associated with free unanchored Lys-63 polyubiquitin chain, promotes autophosphorylation and subsequent activation of MAP3K7. Dephosphorylation


at Thr-187 by PP2A and PPP6C leads to inactivation.

Ubiquitinated, leading to proteasomal degradation (By similarity). Requires 'Lys-63'-linked

polyubiquitination for autophosphorylation and subsequent activation. 'Lys-63'-linked

ubiquitination does not lead to proteasomal degradation. Deubiquitinated by CYLD, a protease that selectively cleaves 'Lys-63'-linked ubiquitin chains. Deubiquitinated by Y.enterocolitica YopP.


## 图片



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-TAK1 antibody [EPR5984] (ab199473)

ab199473 staining TAK1 in wild-type HAP1 cells (top panel) and MAP3K7 knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab199473 at 1/250 dilution and <a href="mailto:ab195887">ab195887</a> at 1/250 dilution (shown in green) overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.


Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-TAK1 antibody [EPR5984] (ab199473)

ab199473 staining TAK1 in differentiated Neuro2a cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab199473 at a 1/100 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor<sup>®</sup> 488), at a 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue). Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in differentiated Neuro2a cells fixed with 4% formaldehyde (10 min).




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-TAK1 antibody [EPR5984] (ab199473)

ab199473 staining TAK1 in MCF7 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab199473 at a 1/100 dilution (shown in red) and <a href="mailto:ab195887">ab195887</a>, Mouse monoclonal to alpha Tubulin (Alexa Fluor<sup>®</sup> 488), at a 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in MCF7 cells fixed with 4% formaldehyde (10 min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors