


### Anti-AMPK alpha 1 antibody [Y365] ab32047

敲除验证
重组
RabMAb

★★★★★
[6 Abreviews](#)
[165 References](#)
[12 图像](#)

#### 概述

产品名称	Anti-AMPK alpha 1抗体[Y365]
描述	兔单克隆抗体[Y365] to AMPK alpha 1
宿主	Rabbit
特异性	This antibody is specific for human AMPK alpha 1. This antibody shows low affinity on mouse and rat samples.
经测试应用	<b>适用于:</b> ICC/IF, Flow Cyt (Intra), WB, IHC-P, IP
种属反应性	<b>与反应:</b> Mouse, Rat, Human <b>预测可用于:</b> African green monkey 
免疫原	Synthetic peptide within Human AMPK alpha 1 aa 500 to the C-terminus (C terminal). The exact sequence is proprietary. Database link: <a href="#">Q13131</a>
阳性对照	WB: HeLa, HepG2, C6, NIH/3T3 and MCF-7 cell lysate. Mouse liver, brain, retina, and skeletal muscle tissue lysates, RAW 264.7 cell lysate, Neuro2A cell lysate. IHC-P: Human cervical carcinoma and lung carcinoma tissues. ICC/IF: MCF-7 cells. Flow Cyt (intra): HeLa cells. IP: HeLa whole cell lysate ( <a href="#">ab150035</a> ).
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> .  Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> .

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20

	Preservative: 0.01% Sodium azide
	Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	Y365
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab32047于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/250.
Flow Cyt (Intra)		1/100 - 1/150. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (5)	1/1000 - 1/5000. Predicted molecular weight: 63 kDa. This antibody shows low affinity on mouse and rat samples.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See <b>IHC antigen retrieval protocols</b> . Mouse and rat species are recommended by WB application.
IP		1/40 - 1/50.

靶标

功能	Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit.
序列相似性	Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily. Contains 1 protein kinase domain.

图片



Western blot - Anti-AMPK alpha 1 antibody [Y365]  
(ab32047)

**All lanes :** Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/1000 dilution

**Lane 1 :** Wild-type RAW 264.7 cell lysate

**Lane 2 :** PRKAA1 knockout RAW 264.7 cell lysate

**Lane 3 :** Mouse Liver cell lysate

**Lane 4 :** Neuro2A cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 63 kDa

**Observed band size:** 64 kDa

False colour image of Western blot: Anti-AMPK alpha 1 antibody [Y365] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab32047 was shown to bind specifically to AMPK alpha 1. A band was observed at 64 kDa in wild-type RAW 264.7 cell lysates with no signal observed at this size in PRKAA1 knockout cell line [ab280055](#) (knockout cell lysate [ab280114](#)). To generate this image, wild-type and PRKAA1 knockout RAW 264.7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween<sup>®</sup> 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L (IRDye<sup>®</sup> 800CW) preabsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye<sup>®</sup> 680RD) preabsorbed ([ab216776](#)) at 1/20000 dilution.



Western blot - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

**Lane 1:** Wild-type HAP1 cell lysate (20 µg)


**Lane 2:** AMPK alpha knockout HAP1 cell lysate (20 µg)

**Lane 3:** MCF7 cell lysate (20 µg)

**Lane 4:** HeLa cell lysate (20 µg)


**Lanes 1 - 4:** Merged signal (red and green). Green - ab32047 observed at 63 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab32047 was shown to specifically react with AMPK alpha in wild-type HAP1 cells. No band was observed when AMPK alpha knockout samples were examined. Wild-type and AMPK alpha knockout samples were subjected to SDS-PAGE. ab32047 and **ab8245** (loading control to GAPDH) were diluted 1/5000 and 1/2000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1/10,000 dilution for 1 hr at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling AMPK alpha 1 with purified ab32047 at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9 before commencing with IHC staining protocol. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.




Immunocytochemistry/ Immunofluorescence - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Immunocytochemistry/Immunofluorescence analysis of MCF7 cells labelling AMPK alpha 1 with purified ab32047 at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.


Control 1: primary antibody (1/250) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).



Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Intracellular Flow Cytometry analysis of HeLa cells labelling AMPK alpha 1 with purified ab32047 at 1/150 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.



ab32047 (purified) at 1/40 immunoprecipitating AMPK alpha 1 in HeLa whole cell lysate.

Lane 1 (input): HeLa whole cell lysate (10µg)


Lane 2 (+): ab32047 + HeLa whole cell lysate (10µg).

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab32047 in HeLa whole cell lysate.

For western blotting, [ab131366](#) VeriBlot for IP Detection Reagent (HRP) was used for detection (1/1500).

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



**All lanes** : Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/500 dilution

**Lane 1** : HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate

**Lane 2** : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

**Lane 3** : Mouse liver tissue lysate

**Lane 4** : Mouse brain tissue lysate

**Lane 5** : Mouse kidney tissue lysate

**Lane 6** : Mouse retina tissue lysate

**Lane 7** : Mouse skeletal muscle tissue lysate

Lysates/proteins at 20 µg per lane.

## Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 63 kDa


**Observed band size:** 63 kDa

**Additional bands at:** 40 kDa (possible non-specific binding)

**Exposure time:** 3 minutes

Blocking/Diluting buffer and concentration: 5% NFDM/TBST.

This antibody shows low affinity on mouse samples.



**All lanes :** Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/20000 dilution

**Lane 1 :** MCF7 (Human breast adenocarcinoma epithelial cell) whole cell lysate

**Lane 2 :** HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

**Lane 3 :** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

**Lane 4 :** K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

**Lane 5 :** C6 (Rat glial tumor glial cell) whole cell lysate

**Lane 6 :** Neuro-2a (Mouse neuroblastoma neuroblast) whole cell lysate

**Lane 7 :** NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

Lysates/proteins at 20 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


**Predicted band size:** 63 kDa

**Observed band size:** 63 kDa

**Exposure time:** 180 seconds

**Blocking and dilution buffer and concentration:** 5% NFDM/TBST.

This antibody shows low affinity on mouse and rat samples.



**All lanes :** Anti-AMPK alpha 1 antibody [Y365] (ab32047) at 1/2000 dilution (purified)

**Lane 1 :** C6 cell lysate

**Lane 2 :** NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary


**All lanes :** Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

**Predicted band size:** 63 kDa

**Observed band size:** 63 kDa


Blocking and dilution buffer: 5% NFDM/TBST.

This antibody shows low affinity on mouse and rat samples.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labelling AMPK alpha 1 with unpurified ab32047 at a dilution of 1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





Flow Cytometry (Intracellular) - Anti-AMPK alpha 1 antibody [Y365] (ab32047)

Overlay histogram showing HeLa cells stained with unpurified ab32047 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified ab32047, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-AMPK alpha 1 antibody [Y365] (ab32047)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors