

# Anti-Amyloid Fibril antibody [mOC87] - Conformation-Specific ab201062

重组 RabMAb

★★★★★ [2 Abreviews](#) [5 References](#) [5 图像](#)

### 概述

产品名称	Anti-Amyloid Fibril抗体[mOC87] - Conformation-Specific
描述	兔单克隆抗体[mOC87] to Amyloid Fibril - Conformation-Specific
宿主	Rabbit
经测试应用	适用于: Dot blot, IHC-P, IHC-FrFI
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide corresponding to Human Amyloid Fibril. Amyloid beta 1-42 fibrils were used as the immunogen. Database link: <a href="#">P05067</a>
阳性对照	beta Amyloid (A $\beta$ ) 1-40; beta Amyloid (A $\beta$ ) 1-42. IHC-P: FFPE human Alzheimer hippocampus tissue sections.
常规说明	<p>This antibody was developed as part of a collaboration between Abcam and Professor Charles Glabe, UC Irvine.</p> <p>ab201062 recognizes a generic epitope of amyloid fibrils and oligomers that is independent of linear sequence (<a href="#">Hatami et al. 2014</a>). Its reactivity with A<math>\beta</math> monomer and oligomers is decreased or eliminated upon thermal denaturation at 100°C of A<math>\beta</math> in SDS sample buffer on western blots (<a href="#">Hatami et al. 2014</a>).</p> <p>For further information on the immunogen, please refer to <a href="#">Hatami et al. 2014</a> and <a href="#">Kayed et al. 2007</a>.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	mOC87
同种型	IgG

## 应用

### The Abpromise guarantee

**Abpromise™** 承诺保证使用ab201062于以下的经测试应用


“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Dot blot		1/8000.
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
IHC-FrFI		Use at an assay dependent concentration.

## 靶标

细胞定位 Membrane.


## 图片



Immunohistochemistry - Free Floating - Anti-Amyloid Fibril antibody [mOC87] - Conformation-Specific (ab201062)

Image courtesy of Professor Charles Glabe, UC Irvine

Immunohistochemical staining of human brain tissue from a patient with a diagnosis of Alzheimers disease, male, 81 years, 5 hour post mortem index, tangle stage 5, plaque stage B, mini mental status exam score 12. Sections were cut using a vibratome. No antigen retrieval was performed. Free floating sections were stained using ab201062 at a dilution of 50 ng/mL. The secondary antibody used was a biotinylated goat anti-rabbit at a dilution of 1/225, which was blocked with normal goat serum. The sample was visualized using ABC solution (1 hour incubation) followed by 1-4 minutes of DAB. The sample was mounted and allowed to dry overnight, followed by dehydration in increasingly concentrated ethanol solutions.



Dot Blot - Anti-Amyloid Fibril antibody [mOC87] - Conformation-Specific (ab201062)

Dot blot analysis of beta Amyloid 1-42 labeled with ab201062 at 1/8000 dilution.

Lane 1: beta Amyloid (A $\beta$ ) 1-40.

Lane 2: beta Amyloid (A $\beta$ ) 1-42.

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/5000 dilution was used as secondary antibody.

Blocking and diluting buffer: 5% NFDm/TBST.

Exposure time: 30 seconds.


Antibody reactivity was assessed using a dot blot, which is a non-quantitative method that maintains the native conformation of beta Amyloid. Beta Amyloid 1-40 and 1-42 peptides underwent the following aggregation conditions before being spotted onto a nitrocellulose membrane and detected using ab201062:

**Monomers:** 0.3 mg of beta Amyloid peptide was dissolved in 30  $\mu$ l 100 mM NaOH and incubated at room temperature for 10 minutes. It was then diluted with 970  $\mu$ l of 1% SDS and boiled for five minutes.

**Oligomers:** 0.3 mg of beta Amyloid peptide was dissolved in 30  $\mu$ l 100 mM NaOH and incubated at room temperature for 10 minutes. It was then diluted with 970  $\mu$ l of 10 mM phosphate buffer pH 7.4 containing 0.02% sodium azide and incubated at room temperature for four days.

**Fibrils:** 0.3 mg of beta Amyloid peptide was dissolved in 1 ml 50% hexafluoroisopropanol (HFIP) with 0.02% sodium azide. It was then stirred constantly for nine days; the first seven with a cap on and the final two with the cap removed to allow evaporation of the HFIP. Fibrils were then sedimented at 20,000 rpm in a microcentrifuge for 20 minutes and resuspended in 1 ml of PBS + 0.02% sodium

azide.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Amyloid Fibril antibody [mOC87] - Conformation-Specific (ab201062)

IHC image of beta Amyloid staining in human Alzheimer hippocampus formalin fixed paraffin embedded tissue section\*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab201062, 1µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

\*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Dot Blot - Anti-Amyloid Fibril antibody [mOC87] - Conformation-Specific (ab201062)

Negative control (secondary ab only):

Lane 1: beta Amyloid (A $\beta$ ) 1-40.

Lane 2: beta Amyloid (A $\beta$ ) 1-42.

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated ([ab97051](#)) at 1/5000 dilution was used as secondary antibody.

Blocking and diluting buffer: 5% NFD/MTBST.

Exposure time: 30 seconds.

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Amyloid Fibril antibody [mOC87] -  
Conformation-Specific (ab201062)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors