abcam

Product datasheet

APC Anti-CD4 antibody [MEM-241] ab18280

1 References 2 图像

概述

产**品名称** APC Anti-CD4**抗体**[MEM-241]

描述 APC小鼠单克隆抗体[MEM-241] to CD4

宿主 Mouse

偶联物 APC. Ex: 645nm, Em: 660nm

经测试应用适用于: Flow Cyt种属反应性与反应: Human

免疫原 Fusion protein (Human).

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C.

存储溶液 pH: 7.40

Preservative: 0.097% Sodium azide

Constituents: PBS, BSA

纯**度** Protein A purified

 克隆
 单克隆

 克隆编号
 MEM-241

 同种型
 IgG1

应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		1/20. ab37391 - Mouse monoclonal lgG1, is suitable for use as an isotype control with this antibody.

靶标

功能 Accessory protein for MHC class-II antigen/T-cell receptor interaction. May regulate T-cell

activation. Induces the aggregation of lipid rafts.

序列相似性 Contains 3 lg-like C2-type (immunoglobulin-like) domains.

Contains 1 lg-like V-type (immunoglobulin-like) domain.

翻译后修饰 Palmitoylation and association with LCK contribute to the enrichment of CD4 in lipid rafts.


Cell membrane. Localizes to lipid rafts. Removed from plasma membrane by HIV-1 Nef protein that increases clathrin-dependent endocytosis of this antigen to target it to lysosomal degradation.

Cell surface expression is also down-modulated by HIV-1 Envelope polyprotein gp160 that

interacts with, and sequesters CD4 in the endoplasmic reticulum.


图片

细胞定位



Flow Cytometry - APC Anti-CD4 antibody [MEM-241] (ab18280)

Surface staining of human peripheral blood cells with anti-human CD4 (MEM-241) APC.



Overlay histogram showing peripheral blood lymphocytes stained with ab18280 (red line). The cells were incubated with the antibody (ab18280, 1/20 dilution) for 30 min at 4°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (ab91353, 2µg/1x10⁶ cells) with secondary antibody DyLight® 649 goat anti-mouse IgG (H&L) (ab96882) at 1/200 dilution for 30 min at 4°C. Acquisition of >5,000 events was performed gating on peripheral blood lymphocytes.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors