

### APC Anti-CD69 antibody [H1.2F3] ab95611

#### 1 图像

#### 概述

产品名称	APC Anti-CD69抗体[H1.2F3]
描述	APC亚美尼亚仓鼠单克隆抗体[H1.2F3] to CD69
宿主	Armenian hamster
偶联物	APC. Ex: 645nm, Em: 660nm
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Mouse
免疫原	Tissue, cells or virus corresponding to Mouse CD69. Murine dendritic epidermal T cell line Y245.
阳性对照	1-day ConA-stimulated BALB/c splenocytes; stimulated Mouse splenic cell cultures
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 7.20 Preservative: 0.09% Sodium azide
纯度	Protein G purified
克隆	单克隆
克隆编号	H1.2F3
同种型	IgG

#### 应用

## The Abpromise guarantee

**Abpromise™** 承诺保证使用ab95611于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use 0.125-0.25µg for 10 <sup>5-8</sup> cells. Staining the cell sample in a final volume of 100 µl is recommended.

## 靶标

功能	Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets.
组织特异性	Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets.
序列相似性	Contains 1 C-type lectin domain.
发展阶段	Earliest inducible cell surface glycoprotein acquired during lymphoid activation.
翻译后修饰	Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes.
细胞定位	Membrane.

## 图片



Staining of 1-day ConA-stimulated BALB/c splenocytes with 0.125 µg of Armenian Hamster IgG Isotype Control APC (blue histogram) or 0.125 µg of ab95611 (purple histogram). Total viable cells were used for analysis.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors