

### Anti-Argonaute-2 antibody [EPR10411] ab186733

敲除验证
重组
RabMAb

163 References 9 图像

#### 概述

产品名称	Anti-Argonaute-2抗体[EPR10411]
描述	兔单克隆抗体[EPR10411] to Argonaute-2
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), ICC/IF, IP, WB, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	HeLa, MCF7, HepG2, C6 and RAW 264.7 cell lysates; Human cervix carcinoma and Mouse kidney tissues; HeLa and MCF7 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol, 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR10411
同种型	IgG

## 应用

### The Abpromise guarantee

**Abpromise™** 承诺保证使用ab186733于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/60. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		1/100 - 1/250.
IP		1/30 - 1/50.
WB		1/1000 - 1/2000. Detects a band of approximately 97 kDa (predicted molecular weight: 97 kDa).
IHC-P		1/100. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

## 靶标

### 功能

Required for RNA-mediated gene silencing (RNAi) by the RNA-induced silencing complex (RISC). The 'minimal RISC' appears to include EIF2C2/AGO2 bound to a short guide RNA such as a microRNA (miRNA) or short interfering RNA (siRNA). These guide RNAs direct RISC to complementary mRNAs that are targets for RISC-mediated gene silencing. The precise mechanism of gene silencing depends on the degree of complementarity between the miRNA or siRNA and its target. Binding of RISC to a perfectly complementary mRNA generally results in silencing due to endonucleolytic cleavage of the mRNA specifically by EIF2C2/AGO2. Binding of RISC to a partially complementary mRNA results in silencing through inhibition of translation, and this is independent of endonuclease activity. May inhibit translation initiation by binding to the 7-methylguanosine cap, thereby preventing the recruitment of the translation initiation factor eIF4-E. May also inhibit translation initiation via interaction with EIF6, which itself binds to the 60S ribosomal subunit and prevents its association with the 40S ribosomal subunit. The inhibition of translational initiation leads to the accumulation of the affected mRNA in cytoplasmic processing bodies (P-bodies), where mRNA degradation may subsequently occur. In some cases RISC-mediated translational repression is also observed for miRNAs that perfectly match the 3' untranslated region (3'-UTR). Can also upregulate the translation of specific mRNAs under certain growth conditions. Binds to the AU element of the 3'-UTR of the TNF (TNF-alpha) mRNA and upregulates translation under conditions of serum starvation. Also required for transcriptional gene silencing (TGS), in which short RNAs known as antigene RNAs or agRNAs direct the transcriptional repression of complementary promoter regions.

### 序列相似性

Belongs to the argonaute family. Ago subfamily.  
Contains 1 PAZ domain.  
Contains 1 Piwi domain.

### 结构域

The Piwi domain may perform RNA cleavage by a mechanism similar to that of RNase H. However while RNase H utilizes a triad of Asp-Asp-Glu (DDE) for metal ion coordination, this protein appears to utilize a triad of Asp-Asp-His (DDH).


翻译后修饰

Hydroxylated. 4-hydroxylation appears to enhance protein stability but is not required for miRNA-binding or endonuclease activity.

细胞定位

Cytoplasm > P-body. Nucleus. Translational repression of mRNAs results in their recruitment to P-bodies. Translocation to the nucleus requires IMP8.

图片



**All lanes :** Anti-Argonaute-2 antibody [EPR10411] (ab186733) at 1/1000 dilution

**Lane 1 :** Wild-type HCT 116 cell lysate

**Lane 2 :** AGO2 knockout HCT 116 cell lysate

**Lane 3 :** U-87 MG cell lysate

**Lane 4 :** MCF7 cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 97 kDa

**Observed band size:** 95 kDa

False colour image of Western blot: Anti-Argonaute-2 antibody [EPR10411] staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] ([ab8245](#)) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab186733 was shown to bind specifically to Argonaute-2. A band was observed at 95 kDa in wild-type HCT 116 cell lysates with no signal observed at this size in AGO2 knockout cell line. To generate this image, wild-type and AGO2 knockout HCT 116 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% milk in TBS-0.1% Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4°C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.



Western blot - Anti-Argonaute-2 antibody  
[EPR10411] (ab186733)

**All lanes** : Anti-Argonaute-2 antibody [EPR10411] (ab186733) at 1/1000 dilution

**Lane 1** : HeLa cell lysate

**Lane 2** : MCF7 cell lysate

**Lane 3** : HepG2 cell lysate


Lysates/proteins at 20 µg per lane.

#### Secondary

**All lanes** : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

**Predicted band size:** 97 kDa


**Observed band size:** 97 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Argonaute-2 antibody  
[EPR10411] (ab186733)


Immunohistochemical analysis of paraffin-embedded Human cervix carcinoma tissue labeling Ago2 / eIF2C2 with ab186733 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Argonaute-2 antibody [EPR10411] (ab186733)


Immunofluorescent analysis of 4% paraformaldehyde-fixed HeLa cells labeling Ago2 / eIF2C2 with ab186733 at 1/250 dilution followed by Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody at 1/200 dilution. Counter stained with Dapi.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Argonaute-2 antibody [EPR10411] (ab186733)

Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling Ago2 / eIF2C2 with ab186733 at 1/100 dilution followed by prediluted HRP Polymer for Rabbit IgG. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Western blot - Anti-Argonaute-2 antibody [EPR10411] (ab186733)

**All lanes** : Anti-Argonaute-2 antibody [EPR10411] (ab186733) at 1/1000 dilution

**Lane 1** : C6 cell lysate

**Lane 2** : RAW 264.7 cell lysate


Lysates/proteins at 10 µg per lane.

**Secondary**

**All lanes** : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution


**Predicted band size:** 97 kDa

**Observed band size:** 97 kDa



Flow Cytometry (Intracellular) - Anti-Argonaute-2 antibody [EPR10411] (ab186733)

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed MCF7 cells labeling Ago2 / eIF2C2 with ab186733 at 1/60 dilution (red) compared to a Rabbit monoclonal IgG isotype control (green), followed by Goat anti rabbit IgG (FITC) secondary antibody at 1/150 dilution.




Immunoprecipitation - Anti-Argonaute-2 antibody [EPR10411] (ab186733)

Western blot analysis of Ago2 / eIF2C2 in MCF7 cell lysate immunoprecipitated with ab186733 at 1/50 dilution.

Secondary antibody: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution.

### Why choose a recombinant antibody?



Anti-Argonaute-2 antibody [EPR10411] (ab186733)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors