abcam

Product datasheet

Anti-ARTS1/ERAP1 antibody [EPR6069] - BSA and Azide free ab232466



1 References 3 图像

概述

产品名称 Anti-ARTS1/ERAP1抗体[EPR6069] - BSA and Azide free

宿主 Rabbit

特异性 The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for

mouse and rat.

种属反应性 与反应: Mouse, Rat, Human

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 IHC-P: Human bladder carcinoma tissue.

常规说明 ab232466 is the carrier-free version of ab124669.

Our <u>carrier-free</u> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C. Do Not Freeze.

存储溶液 pH: 7.2

Constituent: PBS

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR6069

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab232466于以下的经测试应用

"应用说明"部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use at an assay dependent concentration. Predicted molecular weight: 107 kDa.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. We do not guarantee IHC-P for mouse and rat.

靶标

功能 Aminopeptidase that plays a central role in peptide trimming, a step required for the generation of

most HLA class I-binding peptides. Peptide trimming is essential to customize longer precursor peptides to fit them to the correct length required for presentation on MHC class I molecules. Strongly prefers substrates 9-16 residues long. Rapidly degrades 13-mer to a 9-mer and then stops. Preferentially hydrolyzes the residue Leu and peptides with a hydrophobic C-terminus, while it has weak activity toward peptides with charged C-terminus. May play a role in the inactivation of peptide hormones. May be involved in the regulation of blood pressure through the


inactivation of angiotensin II and/or the generation of bradykinin in the kidney.

组织特异性 Ubiquitous.

序列相似性 Belongs to the peptidase M1 family.

翻译后修饰 N-glycosylated.

细**胞定位** Endoplasmic reticulum membrane.




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ARTS1/ERAP1 antibody [EPR6069] - BSA and Azide free (ab232466)

Unpurified <u>ab124669</u>, at a 1/50 dilution, staining ARTS1/ERAP1 in paraffin embedded Human uterus tissue by Immunohistochemistry.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab124669).


Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ARTS1/ERAP1 antibody
[EPR6069] - BSA and Azide free (ab232466)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human bladder carcinoma tissue sections labeling ARTS1/ERAP1 with Purified ab124669 at 1:1000 dilution. Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (<u>ab124669</u>).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

Azide free (ab232466)

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors