


Anti-Axin 2 antibody [EPR2005(2)] ab109307

重组 RabMAb

★★★★★ **5 Abreviews** **49 References** **9 图像**

概述

产品名称	Anti-Axin 2抗体[EPR2005(2)]
描述	兔单克隆抗体[EPR2005(2)] to Axin 2
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Pig 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: C6, PC-12, NIH/3T3, MCF7, SW480, and PC3 cell lysates. IHC-P: Human lung carcinoma tissue. ICC/IF: LnCap cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2005(2)

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab109307于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★★★★★ (2)	1/150.
IHC-P	★★★★★ (2)	1/150. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
WB	★★★★★ (1)	1/1000 - 1/2000. Detects a band of approximately 95 kDa (predicted molecular weight: 94 kDa).

靶标

功能	Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate the phosphorylation of beta-catenin and APC by GSK3B.
组织特异性	Expressed in brain and lymphoblast.
疾病相关	Defects in AXIN2 are involved in colorectal cancer (CRC) [MIM:114500]. They appear to be specifically associated with defective mismatch repair. Defects in AXIN2 are the cause of oligodontia-colorectal cancer syndrome (ODCRCS) [MIM:608615]. Affected individuals manifest severe tooth agenesis and colorectal cancer or precancerous lesions of variable types.
序列相似性	Contains 1 DIX domain. Contains 1 RGS domain.
结构域	The tankyrase-binding motif (also named TBD) is required for interaction with tankyrase TNKS and TNKS2.
翻译后修饰	Probably phosphorylated by GSK3B and dephosphorylated by PP2A. ADP-ribosylated by tankyrase TNKS and TNKS2. Poly-ADP-ribosylated protein is recognized by RNF146, followed by ubiquitination and subsequent activation of the Wnt signaling pathway. Ubiquitinated by RNF146 when poly-ADP-ribosylated, leading to its degradation and subsequent activation of the Wnt signaling pathway. Deubiquitinated by USP34, deubiquitinated downstream of beta-catenin stabilization step: deubiquitination is important Wnt signaling to positively regulate beta-catenin (CTNBB1)-mediated transcription.
细胞定位	Cytoplasm.

图片



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (unpurified)

Lane 1 : SW480 cell lysate

Lane 2 : C6 cell lysate

Lane 3 : PC-12 cell lysate

Lane 4 : NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 94 kDa

Observed band size: 94 kDa


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with unpurified ab109307 at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

This image is courtesy of an anonymous Abreview

Unpurified ab109307 staining Axin 2 in 293T cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized with 1% Triton X-100 and blocked with 3% BSA for 1 hour at room temperature. Samples were incubated with primary antibody (1/50 in PBS + 3% BSA) for 16 hours. An Alexa Fluor® 488-conjugated donkey anti-rabbit IgG polyclonal (1/500) was used as the secondary antibody.



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (purified)

Lane 1 : SW480 cell lysate

Lane 2 : C6 cell lysate

Lane 3 : PC-12 cell lysate

Lane 4 : NIH/3T3 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 94 kDa

Observed band size: 94 kDa


Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with purified ab109307 at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)



Western blot - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

All lanes : Anti-Axin 2 antibody [EPR2005(2)] (ab109307) at 1/1000 dilution (unpurified)

Lane 1 : MCF7 cell lysates


Lane 2 : SW480 cell lysates

Lane 3 : PC3 cell lysates

Lysates/proteins at 10 µg per lane.

Predicted band size: 94 kDa


Observed band size: 95 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with unpurified ab109307 at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor® 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, **ab150113**, an Alexa Fluor® 488-conjugated goat anti-mouse IgG (1/500).



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with purified ab109307 at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor® 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, **ab150113**, an Alexa Fluor® 488-conjugated goat anti-mouse IgG (1/500).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Axin 2 antibody [EPR2005(2)] (ab109307)

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors