


# Anti-Axin 2 antibody [EPR2005(2)] - BSA and Azide free ab192230

**重组** RabMAb

## 6 图像

### 概述

产品名称	Anti-Axin 2抗体[EPR2005(2)] - BSA and Azide free
描述	兔单克隆抗体[EPR2005(2)] to Axin 2 - BSA and Azide free
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Pig 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	MCF7, SW480, and PC3 cell lysates
常规说明	<p>ab192230 is the carrier-free version of <a href="#">ab109307</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
纯化说明	Protein-A purification via MabSelect SuRe
克隆	单克隆
克隆编号	EPR2005(2)
同种型	IgG

## 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab192230于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 95 kDa (predicted molecular weight: 94 kDa).

## 靶标


功能	Inhibitor of the Wnt signaling pathway. Down-regulates beta-catenin. Probably facilitate the phosphorylation of beta-catenin and APC by GSK3B.
组织特异性	Expressed in brain and lymphoblast.
疾病相关	Defects in AXIN2 are involved in colorectal cancer (CRC) [MIM:114500]. They appear to be specifically associated with defective mismatch repair. Defects in AXIN2 are the cause of oligodontia-colorectal cancer syndrome (ODCRCS) [MIM:608615]. Affected individuals manifest severe tooth agenesis and colorectal cancer or precancerous lesions of variable types.
序列相似性	Contains 1 DIX domain. Contains 1 RGS domain.
结构域	The tankyrase-binding motif (also named TBD) is required for interaction with tankyrase TNKS and TNKS2.
翻译后修饰	Probably phosphorylated by GSK3B and dephosphorylated by PP2A.

ADP-ribosylated by tankyrase TNKS and TNKS2. Poly-ADP-ribosylated protein is recognized by RNF146, followed by ubiquitination and subsequent activation of the Wnt signaling pathway. Ubiquitinated by RNF146 when poly-ADP-ribosylated, leading to its degradation and subsequent activation of the Wnt signaling pathway. Deubiquitinated by USP34, deubiquitinated downstream of beta-catenin stabilization step: deubiquitination is important Wnt signaling to positively regulate beta-catenin (CTNBB1)-mediated transcription.

## 细胞定位

Cytoplasm.


## 图片



Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with purified **ab109307** at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor® 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, **ab150113**, an Alexa Fluor® 488-conjugated goat anti-mouse IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab109307**).



Immunocytochemistry/Immunofluorescence analysis of LnCap cells labelling Axin 2 with unpurified **ab109307** at 1/150. Cells were fixed with 4% paraformaldehyde. An Alexa Fluor® 555-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/150) and secondary antibody, **ab150113**, an Alexa Fluor® 488-conjugated goat anti-mouse IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab109307**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] - BSA and Azide free (ab192230)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with purified **ab109307** at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab109307**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Axin 2 antibody [EPR2005(2)] - BSA and Azide free (ab192230)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue labelling Axin 2 with unpurified **ab109307** at 1/150. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. A prediluted HRP-polymer conjugated anti-rabbit IgG was used as the secondary antibody. Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab109307**).



Immunocytochemistry/ Immunofluorescence - Anti-Axin 2 antibody [EPR2005(2)] - BSA and Azide free (ab192230)

This image is courtesy of an anonymous Abreview.

Unpurified **ab109307** staining Axin 2 in 293T cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde, permeabilized with 1% Triton X-100 and blocked with 3% BSA for 1 hour at room temperature. Samples were incubated with primary antibody (1/50 in PBS + 3% BSA) for 16 hours. An Alexa Fluor® 488-conjugated donkey anti-rabbit IgG polyclonal (1/500) was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab109307**).

#### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Axin 2 antibody [EPR2005(2)] - BSA and Azide free (ab192230)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors