

Anti-Axl antibody [MM0098-2N33] ab89224

★★★★★ [4 Abreviews](#) [6 References](#) [1 图像](#)

概述

产品名称	Anti-Axl抗体[MM0098-2N33]
描述	小鼠单克隆抗体[MM0098-2N33] to Axl
宿主	Mouse
经测试应用	适用于: ICC/IF
种属反应性	与反应: African green monkey
免疫原	Human recombinant Axl (extracellular region).
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Constituent: PBS
纯度	Protein G purified
纯化说明	The IgG fraction of culture supernatant was purified by Protein G affinity chromatography.
克隆	单克隆
克隆编号	MM0098-2N33
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用ab89224于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★★★★★ (1)	Use at an assay dependent concentration.

靶标

功能	May function as a signal transducer between specific cell types of mesodermal origin. In case of filovirus infection, seems to function as a cell entry factor.
组织特异性	Highly expressed in metastatic colon tumors. Expressed in primary colon tumors. Weakly expressed in normal colon tissue.
序列相似性	Belongs to the protein kinase superfamily. Tyr protein kinase family. AXL/UFO subfamily. Contains 2 fibronectin type-III domains. Contains 2 Ig-like C2-type (immunoglobulin-like) domains. Contains 1 protein kinase domain.
细胞定位	Membrane.

图片



ab89224 staining Axl in african green monkey CV1 cells by Immunocytochemistry/ Immunofluorescence. The cells were fixed in 4% paraformaldehyde and then blocked using 5% serum for 1 hour at 22°C. Samples were then incubated with primary antibody at 1/100 for 16 hours at 4°C. The secondary antibody used was a donkey anti-mouse IgG conjugated to Alexa Fluor® 546 (red) used at a 1/400 dilution. DAPI was used to counterstain nuclei.

Immunocytochemistry/ Immunofluorescence - Anti-Axl antibody [MM0098-2N33] (ab89224)
Image courtesy of Nir Drayman by Abreview.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors