

Anti-Bcl-XL antibody [E18] - BSA and Azide free ab199099

重组 RabMAb

17 References **11 图像**


概述

产品名称	Anti-Bcl-XL抗体[E18] - BSA and Azide free
描述	兔单克隆抗体[E18] to Bcl-XL - BSA and Azide free
宿主	Rabbit
特异性	This antibody should recognize Bcl-XL, Bcl-xS and Bcl-x(beta) as the immunogen sequence is common to them. The antibody does not cross-react with other Bcl-2 family members.
经测试应用	适用于: Flow Cyt (Intra), ICC/IF, IP, IHC-P, WB
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Pig 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Jurkat whole cell lysate (ab7899) can be used as a positive control in WB.
常规说明	<p>ab199099 is the carrier-free version of ab32370.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
解离常数 (K_D)	$K_D = 6.50 \times 10^{-11}$ M



存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	E18
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab199099 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols .
WB		Use at an assay dependent concentration. Predicted molecular weight: 26 kDa. Please check the parent abID, ab32370 , for more information on dilutions.

标签

功能	Potent inhibitor of cell death. Inhibits activation of caspases (By similarity). Appears to regulate cell death by blocking the voltage-dependent anion channel (VDAC) by binding to it and preventing the release of the caspase activator, CYC1, from the mitochondrial membrane. Isoform Bcl-X(S) promotes apoptosis.
组织特异性	Bcl-X(S) is expressed at high levels in cells that undergo a high rate of turnover, such as developing lymphocytes. In contrast, Bcl-X(L) is found in tissues containing long-lived postmitotic cells, such as adult brain.
序列相似性	Belongs to the Bcl-2 family.
结构域	The BH4 motif is required for anti-apoptotic activity. The BH1 and BH2 motifs are required for both heterodimerization with other Bcl-2 family members and for repression of cell death.
翻译后修饰	Proteolytically cleaved by caspases during apoptosis. The cleaved protein, lacking the BH4 motif, has pro-apoptotic activity.
细胞定位	Mitochondrion membrane. Nucleus membrane. Mitochondrial membranes and perinuclear envelope.

图片



Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099) + C6
(rat glioma) whole cell lysate at 10 µg

Secondary


Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#))

Predicted band size: 26 kDa

Exposure time: 30 seconds

Blocking buffer and concentration: 5% NFDM/TBST

Diluting buffer and concentration: 5% NFDM/TBST




Immunocytochemistry/ Immunofluorescence - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling Bcl-XL with purified **ab32370** at 1/500. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, a goat anti-rabbit Alexa Fluor® 488 (IgG; 1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/500) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human endometrium tissue labelling Bcl-XL with purified **ab32370** at 1/2000. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit HRP (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



Flow Cytometry (Intracellular) - Anti-Bcl-XL antibody
[E18] - BSA and Azide free (ab199099)

Intracellular Flow Cytometry analysis of Jurkat cells labelling Bcl-XL with purified **ab32370** at 1/20 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).




Immunoprecipitation - Anti-Bcl-XL antibody [E18] -
BSA and Azide free (ab199099)

ab32370 (purified) at 1/30 immunoprecipitating Bcl-XL in Jurkat cell lysate (Lane 1). Lane 2 - PBS. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the secondary antibody (1/1500).

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Bcl-XL antibody [E18] -
BSA and Azide free (ab199099)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human prostate carcinoma tissue labelling Bcl-XL with unpurified **ab32370** at 1/50.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



Immunocytochemistry/ Immunofluorescence - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

ICC/IF image of unpurified **ab32370** stained HepG2 cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (**ab32370**, 1/100) overnight at +4°C. The secondary antibody (green) was **ab96899**, goat **anti-rabbit DyLight® 488** (IgG; H+L) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

Image from Medic S & Ziman M PLoS One. 2010 Apr 22;5(4):e9977. Fig 5.; doi:10.1371/journal.pone.0009977; April 22 2010 PLoS ONE 5(4): e9977.

Immunohistochemistry of human primary melanoma, staining Bcl-XL (red) with unpurified **ab32370**.

Antigen retrieval was performed in EDTA/Tris buffer (pH 8) before being blocked with 10%NGS for one hour at room temperature. Samples were incubated with primary antibody (1/50) at room temperature for one hour. An AlexaFluor®-conjugated anti-rabbit IgG was used as the secondary antibody.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



Flow Cytometry (Intracellular) - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

Overlay histogram showing DU145 cells stained with unpurified **ab32370** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab32370**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was a goat anti-rabbit DyLight® 488 (IgG; H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).



BI-RD Scanning - Anti-Bcl-XL antibody [E18] - BSA and Azide free (ab199099)

Equilibrium disassociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab32370**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Bcl-XL antibody [E18] - BSA and Azide free
(ab199099)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors