

Anti-BDNF antibody [EPR1292] ab108319

重组 RabMAb

★★★★★ **8 Abreviews** **380 References** 12 图像

概述

产品名称	Anti-BDNF抗体[EPR1292]
描述	兔单克隆抗体[EPR1292] to BDNF
宿主	Rabbit
特异性	This product may cross react with the following family members: NGF beta, neurotrophin 3, neurotrophin 4. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
经测试应用	适用于: Flow Cyt (Intra), WB, IHC-P, IHC-Fr, ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. (Peptide available as ab182199)
阳性对照	WB: Human, rat and mouse brain, hippocampus and cerebellum lysates; IHC-P: Human brain tissue, human bladder cancer tissue; ICC/IF: HeLa cells; Flow Cyt (intra): HeLa cells; IHC-Fr: Mouse and Rat cerebrum tissue, Hu cerebral cortex.
常规说明	<p>For BDNF, multiple WB bands are possible and expected. The human protein has 5 isoforms (precursors: 28 – 37 kDa) and can be glycosylated (Uniprot: http://www.uniprot.org/uniprot/P23560). The mature form is expected at ~14 kDa (monomer) and the dimer at ~28 kDa.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

	term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR1292
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab108319于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
Flow Cyt (Intra)		1/30.
WB	★★★★★ (5)	1/1000 - 1/10000. Predicted molecular weight: 15 kDa.Can be blocked with Human BDNF peptide (ab182199) .
IHC-P	★★★★★ (1)	1/500. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat. See IHC antigen retrieval protocols .
IHC-Fr		1/100. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)
ICC/IF	★★★★★ (1)	1/500. For unpurified use at 1/750 dilution.

靶标

功能	During development, promotes the survival and differentiation of selected neuronal populations of the peripheral and central nervous systems. Participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. Major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability.
组织特异性	Brain. Highly expressed in hippocampus, amygdala, cerebral cortex and cerebellum. Also expressed in heart, lung, skeletal muscle, testis, prostate and placenta.
疾病相关	Bulimia nervosa 2 Congenital central hypoventilation syndrome

序列相似性	Belongs to the NGF-beta family.
翻译后修饰	The propeptide is N-glycosylated and glycosulfated. Converted into mature BDNF by plasmin (PLG).
细胞定位	Secreted.

图片



All lanes : Anti-BDNF antibody [EPR1292] (ab108319) at 1/1000 dilution (unpurified)

- Lane 1 :** Human hippocampus lysate
- Lane 2 :** Rat hippocampus lysate
- Lane 3 :** Mouse hippocampus lysate


Lysates/proteins at 20 µg per lane.

Secondary
All lanes : Gt anti Rb IR680 at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 15 kDa
Additional bands at: 15 kDa (possible mature (processed) protein), 28 kDa (possible truncated form), 35 kDa (possible immature (unprocessed)), 45 kDa (possible immature (unprocessed))

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using Licor blocking buffer before being incubated with unpurified ab108319 (1/1000) overnight at 4°C. Ab8245 (mouse anti-GAPDH; 0.05 ug/mL) was included as a loading control. Antibody binding was detected using goat anti-rabbit IgG IR-680 (green) and goat anti-mouse IgG IR800 (red) at a 1:10,000 dilution for 1hr at room temperature and then imaged using the Licor Odyssey CLx



All lanes : Anti-BDNF antibody [EPR1292] (ab108319) at 1/1000 dilution (Purified)

Lane 1 : Human brain lysates with 5% NFDM/TBST

Lane 2 : Mouse brain lysates with 5% NFDM/TBST

Lane 3 : Rat brain lysates with 5% NFDM/TBST

Lane 4 : Human hippocampus lysates with 5% NFDM/TBST

Lane 5 : Mouse hippocampus lysates with 5% NFDM/TBST

Lane 6 : Rat hippocampus lysates with 5% NFDM/TBST

Lane 7 : Human cerebellum lysates with 5% NFDM/TBST

Lane 8 : Mouse cerebellum lysates with 5% NFDM/TBST

Lane 9 : Rat cerebellum lysates with 5% NFDM/TBST


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution


Predicted band size: 15 kDa

Observed band size: 15-45 kDa




IHC image of BDNF staining in a section of frozen normal human cerebral cortex performed on a Leica BOND™ system using the standard protocol. The section was fixed in 10% paraformaldehyde (10 min) prior to staining. The section was incubated with ab108319, 1/200 dilution, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. The inset secondary-only control image is taken from an identical assay without primary antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.




Immunohistochemistry (Frozen sections) - Anti-BDNF antibody [EPR1292] (ab108319)

Immunohistochemistry (Frozen sections) analysis of rat cerebral cortex tissue sections labeling BDNF with Purified ab108319 at 1/100 (2.8 µg/ml). Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. DAPI was used as a counterstain.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BDNF antibody [EPR1292] (ab108319)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human bladder cancer tissue sections labeling BDNF with Purified ab108319 at 1:500 dilution (0.56 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0)




Flow Cytometry (Intracellular) - Anti-BDNF antibody
[EPR1292] (ab108319)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling BDNF with purified ab108319 at 1/30 dilution (10µg/ml) (red). Cells were fixed with 80% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).




Western blot - Anti-BDNF antibody [EPR1292]
(ab108319)

Different batches of ab108319 were tested on Mouse brain lysate at 0.3 µg/ml. 15 µg of lysate was loaded in each lane. Bands observed at 14-45 kDa.




Immunohistochemistry (Frozen sections) - Anti-BDNF antibody [EPR1292] (ab108319)

Immunohistochemistry (Frozen sections) analysis of mouse cerebrum tissue sections labeling BDNF with Purified ab108319 at 1/100 (2.8 µg/ml). Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. DAPI was used as a counterstain.



Immunocytochemistry/ Immunofluorescence - Anti-BDNF antibody [EPR1292] (ab108319)


Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling BDNF with Purified ab108319 at 1:500 (0.6 µg/ml). Cells were fixed in 100% Methanol and permeabilized with None. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BDNF antibody [EPR1292] (ab108319)

Immunohistochemical analysis of paraffin-embedded human brain tissue using unpurified ab108319 at 1/100 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-BDNF antibody [EPR1292] (ab108319)

This image is courtesy of an Abreview submitted by Kirk McManus (1894904)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (human cervix adenocarcinoma) cells labelling BDNF with unpurified ab108319 at a dilution of 1/750. Cells were fixed with paraformaldehyde and permeabilized with 0.5% Triton-X100 in PBS. **ab150081** (1/200) was used as the secondary antibody.

The antibody produces a strong, golgi-associated labelling pattern in both PF and MeOH fixed samples.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-BDNF antibody [EPR1292] (ab108319)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors