# abcam

## Product datasheet

## Biotin Anti-CD11c antibody [N418] ab95765

8 References 1 图像

概述

产**品名称** 生物素Anti-CD11c抗体[N418]

描述 生物素亚美尼亚仓鼠单克隆抗体[N418] to CD11c

宿主 Armenian hamster

**偶联物** Biotin

经测试应用 适用于: Flow Cyt 种属反应性 与反应: Mouse

免疫原 Tissue, cells or virus corresponding to Mouse CD11c. Mouse spleen dendritic cells

阳性对照 Flow Cyt: C57BL/6 splenocytes.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

**存放说明** Shipped at 4°C. Store at +4°C. Do Not Freeze. Store In the Dark.

存储溶液 pH: 7.20

Preservative: 0.09% Sodium azide

Constituents: 0.88% Sodium chloride, 0.1% Gelatin, 0.12% Monobasic dihydrogen sodium

phosphate

纯**度** Affinity purified

 克隆
 单克隆

 克隆编号
 N418

**同种型** lgG

#### The Abpromise guarantee

#### Abpromise™承诺保证使用ab95765于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应 <b>用</b>	Ab评论	说明
Flow Cyt		Use 0.25-0.5 $\mu$ g for 10 <sup>5-8</sup> cells. Final volume of 100 $\mu$ l.

#### 靶标

功能 Integrin alpha-X/beta-2 is a receptor for fibrinogen. It recognizes the sequence G-P-R in

fibrinogen. It mediates cell-cell interaction during inflammatory responses. It is especially

important in monocyte adhesion and chemotaxis.

组织特异性 Predominantly expressed in monocytes and granulocytes.

序列相似性 Belongs to the integrin alpha chain family.


Contains 7 FG-GAP repeats. Contains 1 VWFA domain.

结构域 The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo

protease cleavage.

细胞定位 Membrane.

#### 图片



C57BI/6 splenocytes were stained with APC-conjugated anti-MHC CLass II and 0.03µg ab95765 (right) or 0.03µg biotin-conjugated armenian hamster IgG isotype control (left), followed by streptavidin PE.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

• Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

#### Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors