

### Biotin Anti-CD69 antibody [H1.2F3] ab25008

#### 2 References

#### 概述

---

产品名称	生物素Anti-CD69抗体[H1.2F3]
描述	生物素亚美尼亚仓鼠单克隆抗体[H1.2F3] to CD69
宿主	Armenian hamster
偶联物	Biotin
经测试应用	适用于: IP, Flow Cyt, IHC-Fr
种属反应性	与反应: Mouse
免疫原	Y245 dendritic epidermal T cell line (Mouse)
常规说明	

ab25008 augments PMA-induced T-cell proliferation and induces redirected lysis of Fc receptor-bearing target cells by NK cells.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

#### 性能

---

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Store In the Dark.
存储溶液	pH: 7.4 Preservative: 0.1% Sodium azide Constituent: PBS
纯度	Affinity purified
Primary antibody说明	ab25008 augments PMA-induced T-cell proliferation and induces redirected lysis of Fc receptor-bearing target cells by NK cells.
克隆	单克隆

克隆编号	H1.2F3
同种型	IgG
轻链类型	lambda

## 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab25008于以下的经测试应用

“应用说明”部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration. Acetone fixed tissues.

## 靶标

功能	Involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets.
组织特异性	Expressed on the surface of activated T-cells, B-cells, natural killer cells, neutrophils, eosinophils, epidermal Langerhans cells and platelets.
序列相似性	Contains 1 C-type lectin domain.
发展阶段	Earliest inducible cell surface glycoprotein acquired during lymphoid activation.
翻译后修饰	Constitutive Ser/Thr phosphorylation in both mature thymocytes and activated T-lymphocytes.
细胞定位	Membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors