

Biotin Anti-HBEGF/DTR antibody [4G10] ab66793

3 图像

概述

产品名称	生物素Anti-HBEGF/DTR抗体[4G10]
描述	生物素小鼠单克隆抗体[4G10] to HBEGF/DTR
宿主	Mouse
偶联物	Biotin
经测试应用	适用于: IP, ICC/IF, WB
种属反应性	与反应: Human 不与反应: Mouse
免疫原	Recombinant fragment corresponding to Human HBEGF/DTR. Database link: Q99075
表位	EGF domain
阳性对照	Vero cell extract carrying human HG-EGF expression vector.
常规说明	Biotin/IgG = 7.5

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 6 Constituents: PBS, 50% Glycerol (glycerin, glycerine)
纯度	Protein G purified
纯化说明	Produced in serum-free medium and purified by combination of chromatography.
克隆	单克隆

克隆编号 4G10
同种型 IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab66793于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		Use at an assay dependent dilution.
ICC/IF		Use at an assay dependent dilution.
WB		Use a concentration of 0.2 - 1 µg/ml. Detects a band of approximately 20-28 kDa (predicted molecular weight: 23 kDa).

靶标


功能 May be involved in macrophage-mediated cellular proliferation. It is mitogenic for fibroblasts and smooth muscle but not endothelial cells. It is able to bind EGF receptors with higher affinity than EGF itself and is a far more potent mitogen for smooth muscle cells than EGF. Also acts as a diphtheria toxin receptor.

序列相似性 Contains 1 EGF-like domain.

翻译后修饰 Several N-termini have been identified by direct sequencing. The forms with N-termini 63, 73 and 74 have been tested and found to be biologically active.
O-linked glycan attachment sites were determined by Edman degradation, O-glycanase digest suggests mucin-type glycosylation (done in HB-EGF purified from histiocytic lymphoma cell line U-937).

细胞定位 Cell membrane and Secreted > extracellular space. Mature HB-EGF is released into the extracellular space and probably binds to a receptor.

图片



Western blot - Biotin Anti-HBEGF/DTR antibody [4G10] (ab66793)

All lanes : Biotin Anti-HBEGF/DTR antibody [4G10] (ab66793) at 1 µg/ml

Lane 1 : Vero (African green monkey kidney epithelial cell) whole cell lysate


Lane 2 : Vero cell extract from cells carrying human HBEGF/DTR expression vector

Lane 3 : Vero cell extract from cells carrying mouse HBEGF/DTR expression vector

Predicted band size: 23 kDa


Observed band size: 20-28 kDa

Additional bands at: 41 kDa. We are unsure as to the identity of these extra bands.



Immunocytochemistry/ Immunofluorescence - Biotin Anti-HBEGF/DTR antibody [4G10] (ab66793)

These images shows Vero cells carrying human HBEGF/DTR expression vector (left hand image) and Vero cells carrying mouse HBEGF/DTR expression vector (right hand image). Cells treated with ab66793, fixed with 4% PFA and reacted with Cys3 conjugated 2nd antibody.



Immunoprecipitation - Biotin Anti-HBEGF/DTR antibody [4G10] (ab66793)

All lanes : HBEGF/DTR antibody [4G10](biotin) - Azide free (ab66793) at 1 µg/ml. Lane 1 : Vero cell extract Lane 2 : Vero cell extract from cells carrying human HBEGF/DTR expression vector. Lane 3 : Vero cell extract from cells carrying mouse HBEGF/DTR expression vector Predicted band size : 23 kDa Observed band size : 20-38 kDa Additional bands at : 41 kDa. We are unsure as to the identity of these extra bands. This image shows Immunoprecipitation of HBEGF/DTR followed by Western blotting/SDS PAGE; the cell surface was biotinylated.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors