

Anti-CaMKI antibody [EPR2217Y] ab68234

重组 RabMAb

★★★★★ **4 Abreviews** **13 References** **5 图像**

概述

产品名称	Anti-CaMKI抗体[EPR2217Y]
描述	兔单克隆抗体[EPR2217Y] to CaMKI
宿主	Rabbit
经测试应用	适用于: WB, IHC-P 不适用于: Flow Cyt or IP
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human CaMKI (N terminal). The exact sequence is proprietary.
阳性对照	SH-SY5Y lysate, human, mouse and rat brain lysate and human kidney tissue sections.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.05% BSA, 40% Glycerol (glycerin, glycerine), 59% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2217Y
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab68234于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (3)	1/5000 - 1/10000. Detects a band of approximately 41 kDa (predicted molecular weight: 41 kDa).
IHC-P	★★★★★ (1)	1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

应用说明

Is unsuitable for Flow Cyt or IP.

靶标

功能

Calcium/calmodulin-dependent protein kinase that operates in the calcium-triggered CaMKK-CaMK1 signaling cascade and, upon calcium influx, regulates transcription activators activity, cell cycle, hormone production, cell differentiation, actin filament organization and neurite outgrowth. Recognizes the substrate consensus sequence [MVLIF]-x-R-x(2)-[ST]-x(3)-[MVLIF]. Regulates axonal extension and growth cone motility in hippocampal and cerebellar nerve cells. Upon NMDA receptor-mediated Ca^{2+} elevation, promotes dendritic growth in hippocampal neurons and is essential in synapses for full long-term potentiation (LTP) and ERK2-dependent translational activation. Downstream of NMDA receptors, promotes the formation of spines and synapses in hippocampal neurons by phosphorylating ARHGEF7/BETAPIX on 'Ser-694', which results in the enhancement of ARHGEF7 activity and activation of RAC1. Promotes neuronal differentiation and neurite outgrowth by activation and phosphorylation of MARK2 on 'Ser-91', 'Ser-92', 'Ser-93' and 'Ser-294'. Promotes nuclear export of HDAC5 and binding to 14-3-3 by phosphorylation of 'Ser-259' and 'Ser-498' in the regulation of muscle cell differentiation. Regulates NUMB-mediated endocytosis by phosphorylation of NUMB on 'Ser-276' and 'Ser-295'. Involved in the regulation of basal and estrogen-stimulated migration of medulloblastoma cells through ARHGEF7/BETAPIX phosphorylation (By similarity). Is required for proper activation of cyclin-D1/CDK4 complex during G1 progression in diploid fibroblasts. Plays a role in K(+) and ANG2-mediated regulation of the aldosterone synthase (CYP11B2) to produce aldosterone in the adrenal cortex. Phosphorylates EIF4G3/eIF4GII. In vitro phosphorylates CREB1, ATF1, CFTR, MYL9 and SYN1/synapsin I.

组织特异性

Widely expressed. Expressed in cells of the zona glomerulosa of the adrenal cortex.

序列相似性

Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CaMK subfamily. Contains 1 protein kinase domain.

结构域

The autoinhibitory domain overlaps with the calmodulin binding region and interacts in the inactive folded state with the catalytic domain as a pseudosubstrate.


翻译后修饰

Phosphorylated by CaMKK1 and CaMKK2 on Thr-177.
Polyubiquitinated by the E3 ubiquitin-protein ligase complex SCF(FBXL12), leading to proteasomal degradation.

细胞定位

Cytoplasm. Nucleus. Predominantly cytoplasmic.

图片



Western blot - Anti-CaMKI antibody [EPR2217Y]
(ab68234)

All lanes : Anti-CaMKI antibody [EPR2217Y] (ab68234)

Lane 1 : HEK-293 (Human embryonic kidney epithelial cell) whole cell lysate

Lane 2 : Human brain tissue lysate

Lane 3 : Mouse brain tissue lysate

Lane 4 : NIH/3T3 (Mouse embryonic fibroblast)

Lane 5 : Rat brain tissue lysate

Lane 6 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lysates/proteins at 40 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 41 kDa

Observed band size: 41 kDa

Exposure time: Lane 1 & 3, 4, 5: 5 seconds. Lane 2 & 6:20 seconds

Blocking and diluting buffer: 5% NFDM /TBST



Western blot - Anti-CaMKI antibody [EPR2217Y]
(ab68234)

All lanes : Anti-CaMKI antibody [EPR2217Y] (ab68234) at 1/5000 dilution

Lane 1 : SH-SY5Y lysate

Lane 2 : human brain tissue lysate

Lane 3 : mouse brain tissue lysate

Lane 4 : rat brain tissue lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution


Predicted band size: 41 kDa

Observed band size: 41 kDa



Immunohistochemical analysis of Paraffin-embedded Human adrenal gland tissue labelling CaMKI with ab68234 at 1/500 dilution, followed by ready to use secondary antibody Goat Anti-Rabbit IgG H&L (HRP polymer). Positive staining on the zona glomerulosa of human adrenal glands. The section was incubated with ab68234 at +4°C overnight. Counter stained with Hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CaMKI antibody [EPR2217Y] (ab68234)



ab68234 at 1/100-1/250 dilution staining CaMKI in human kidney tissue sections.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CaMKI antibody [EPR2217Y] (ab68234)

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-CaMKI antibody [EPR2217Y] (ab68234)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors