abcam

Product datasheet

Anti-Cardiac Troponin T antibody [1C11] ab8295

★★★★★ 14 Abreviews 243 References 4 图像

概述

产**品名称** Anti-Cardiac Troponin T抗体[1C11]

小鼠单克隆抗体[1C11] to Cardiac Troponin T

宿主 Mouse

经测试应用 适用于: ICC/IF, IHC-P, Sandwich ELISA

种属反应性 与反应: Mouse, Rat, Dog, Human

免疫原 Other Immunogen Type corresponding to Human Cardiac Troponin T aa 171-190.

Database link: P45379

阳性对照 Natural Human Cardiac Troponin T protein (ab9937) can be used as a positive control in WB.

IHC: Human heart tissue. ICC/IF: ioSkeletal Myocytes - Human iPSC-Derived Skeletal Myocytes

(ab277612).

常规说明 This antibody detects Troponin T in human cardiac muscle. No cross-reaction with skeletal

troponin T, cTnl and TnC.

This product was changed from ascites to tissue culture supernatant on 17th October 2017 and

product received after this date will be from tissue culture supernatant.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

存储溶液 pH: 7.40

Preservative: 0.1% Sodium azide

纯度 Protein A purified **纯化说明** Purified from TCS

1

Primary antibody说明 This antibody detects Troponin T in human cardiac muscle. No cross-reaction with skeletal

troponin T, cTnl and TnC.

 克隆
 单克隆

 克隆编号
 1C11

 骨髓瘤
 Sp2/0

同种型 lgG1

轻链类型 unknown

应用

The Abpromise guarantee Abpromise™承诺保证使用ab8295于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF	★★★★ (6)	Use a concentration of 1 - 5 μg/ml.
IHC-P	**** <u>(2)</u>	Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
Sandwich ELISA		Use at an assay dependent concentration. Can be used as Capture or Detection antibody.

靶标

功能 Troponin T is the tropomyosin-binding subunit of troponin, the thin filament regulatory complex

which confers calcium-sensitivity to striated muscle actomyosin ATPase activity.

组织**特异性** Heart. The fetal heart shows a greater expression in the atrium than in the ventricle, while the adult

heart shows a greater expression in the ventricle than in the atrium. Isoform 6 predominates in normal adult heart. Isoforms 1, 7 and 8 are expressed in fetal heart. Isoform 7 is also expressed in

failing adult heart.

疾病相关 Defects in TNNT2 are the cause of cardiomyopathy familial hypertrophic type 2 (CMH2)

[MIM:115195]. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.

Defects in TNNT2 are the cause of cardiomyopathy dilated type 1D (CMD1D) [MIM:601494].

Dilated cardiomyopathy is a disorder characterized by ventricular dilation and impaired systolic

function, resulting in congestive heart failure and arrhythmia. Patients are at risk of premature

death.


Defects in TNNT2 are the cause of cardiomyopathy familial restrictive type 3 (RCM3) $\,$

[MIM:612422]. Restrictive cardiomyopathy is a heart disorder characterized by impaired filling of the ventricles with reduced diastolic volume, in the presence of normal or near normal wall

thickness and systolic function.

序列相似性 Belongs to the troponin T family.


2



Paraffin-embedded Normal human Heart and iVSD heart tissue (were blocked using 10% FBS for 30 min) stained for Cardiac Troponin T (Red) using ab8295 at 1/200 dilution at room temperature for 2 hours. The slides were then incubated with Fluor® 555-conjugated anti-mouse (Abcam, <u>ab150107</u>; 1:1,000 dilution). The nuclear counterstain was DAPI (Blue).

Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

Ye L et al. Decreased Yes-Associated Protein-1 (YAP1) Expression in Pediatric Hearts with Ventricular Septal Defects. PLoS One 10:e0139712 (2015).




Immunocytochemistry/ Immunofluorescence - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

Immunofluorescence staining of Cardiac Troponin T using ab8295 in ioSkeletal Myocytes - Human iPSC-Derived Skeletal Myocytes (ab277612), which were differentiated for 10 days post induction.

The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% PBS-Tween for 5 mins and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab8295 at 5 µg/mL and ab6046, rabbit polyclonal to beta Tubulin, at 1/1000 dilution. Cells were then incubated with ab150117, Goat Anti-Mouse lgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and ab150088, Goat Anti-Rabbit lgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).


Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown. Gamma is adjusted to 1.5 in all channels.

The antibody ab8295 gave comparable results using MeOH fixation (100%, $5\,\mathrm{min}$).



Sandwich ELISA - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

Calibration curves for sandwich cTnT fluoroimmunoassay with different animal TnTs as antigen.(dark blue) canine, (blue/grey) human, (grey) mouse, (black) rat. Monoclonal antibodies: capture, ab8295 [clone 1C11], 1 μ g/well, detection <u>ab1454</u> [clone 7E7], 200 ng/well. Assay time, 30 min at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Cardiac Troponin T antibody [1C11] (ab8295)

Ab8295 staining human normal heart. Staining is localised to the cytoplasm.

Left panel: with primary antibody at 1 ug/ml. Right panel: isotype control.

Sections were stained using an automated system DAKO Autostainer Plus, at room temperature. Sections were rehydrated and antigen retrieved with the Dako 3-in-1 AR buffer citrate pH 6.0 in a DAKO PT Link. Slides were peroxidase blocked in 3% H2O2 in methanol for 10 minutes. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS), then incubated with primary antibody for 20 minutes, and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors