

Anti-CBL antibody [YE323] - C-terminal ab32027

敲除验证
重组
RabMAb

11 References 7 图像

概述

产品名称	Anti-CBL 抗体[YE323] - C-terminal
描述	兔单克隆抗体[YE323] to CBL - C-terminal
宿主	Rabbit
经测试应用	适用于: WB, Flow Cyt (Intra), ICC/IF 不适用于: IP
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Chicken 
免疫原	Synthetic peptide within Human CBL aa 850 to the C-terminus (C terminal). The exact sequence is proprietary. Database link: P22681
表位	ab32027 reacts with an epitope located in the C terminal region of CBL.
阳性对照	WB: HEK293T, HAP1, Jurkat, THP-1, WEHI-231, F9 and Raji cell lysates; Mouse thymus tissue lysate, Rat testis lysate, Rat thymus lysate. ICC/IF: Jurkat cells. Flow Cyt (intra): Jurkat cells.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

纯度	Protein A purified
克隆	单克隆
克隆编号	YE323
同种型	IgG

应用

The Abpromise guarantee **Abpromise™承诺保证使用ab32027于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 120 kDa (predicted molecular weight: 99 kDa). For unpurified use at 1/5000
Flow Cyt (Intra)		1/30. For unpurified use at 1/100 <u>ab172730</u> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		1/50. For unpurified use at 1/100

应用说明 Is unsuitable for IP.

靶标

功能	Participates in signal transduction in hematopoietic cells. Adapter protein that functions as a negative regulator of many signaling pathways that start from receptors at the cell surface. Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including PDGFA, EGF and CSF1, and terminates signaling.
-----------	--

通路 Protein modification; protein ubiquitination.

疾病相关	<p>Defects in CBL are the cause of Noonan syndrome-like disorder (NSL) [MIM:613563]. NSL is a syndrome characterized by a phenotype reminiscent of Noonan syndrome. Clinical features are highly variable, including facial dysmorphism, short neck, developmental delay, hyperextensible joints and thorax abnormalities with widely spaced nipples. The facial features consist of triangular face with hypertelorism, large low-set ears, ptosis, and flat nasal bridge. Some patients manifest cardiac defects.</p>
------	---

序列相似性	Contains 1 Cbl-PTB (Cbl-type phosphotyrosine-binding) domain. Contains 1 RING-type zinc finger. Contains 1 UBA domain.
-------	--

结构域 The RING-type zinc finger domain mediates binding to an E2 ubiquitin-conjugating enzyme. The N-terminus is composed of the phosphotyrosine binding (PTB) domain, a short linker region and the RING-type zinc finger. The PTB domain, which is also called TKB (tyrosine kinase binding) domain, is composed of three different subdomains: a four-helix bundle (4H), a calcium-binding EF hand and a divergent SH2 domain.


翻译后修饰 Phosphorylated on tyrosine residues by EGFR, SYK, FYN and ZAP70 (By similarity).

Phosphorylated on tyrosine residues by INSR.

细胞定位

Cytoplasm.

图片



All lanes : Anti-CBL antibody [YE323] - C-terminal (ab32027) at 1/1000 dilution

Lane 1 : Raji (Human Burkitt's lymphoma B lymphocyte) whole cell lysate at 15 µg

Lane 2 : Rat testis lysate at 20 µg


Lane 3 : Rat thymus lysate at 20 µg

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/2000 dilution

Predicted band size: 99 kDa

Observed band size: 110 kDa



All lanes : Anti-CBL antibody [YE323] - C-terminal (ab32027) at 1/1000 dilution (unpurified)

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : CBL knockout HEK293T cell lysate

Lane 3 : THP-1 cell lysate

Lane 4 : Raji cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

Predicted band size: 99 kDa

Observed band size: 110 kDa


Lanes 1-4: Merged signal (red and green). Green - ab32027 observed at 110 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

Unpurified ab32027 Anti-CBL antibody [YE323] - C-terminal was shown to specifically react with CBL in wild-type HEK293T cells. Loss of signal was observed when knockout cell line **ab267245** (knockout cell lysate **ab257200**) was used. Wild-type and CBL knockout samples were subjected to SDS-PAGE. ab32027 and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-CBL antibody [YE323] - C-terminal (ab32027)

Immunocytochemistry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling CBL with purified ab32027 at 1/50 dilution (4.26 µg/mL). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/mL). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1/1000 (2 µg/mL) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-CBL antibody [YE323] - C-terminal (ab32027)

Intracellular Flow Cytometry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling CBL with purified ab32027 at 1/30 dilution (10 µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-CBL antibody [YE323] - C-terminal (ab32027)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: CBL knockout HAP1 whole cell lysate (20 µg)

Lane 3: THP1 whole cell lysate (20 µg)

Lane 4: Raji whole cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green). Green - ab32027 observed at 100 kDa. Red - loading control, [ab9484](#), observed at 37 kDa.

Unpurified ab32027 was shown to specifically react with CBL in wild-type HAP1 cells as signal was lost in CBL knockout cells. Wild-type and CBL knockout samples were subjected to SDS-PAGE. ab32027 and [ab9484](#) (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/5000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-CBL antibody [YE323] - C-terminal (ab32027)

All lanes : Anti-CBL antibody [YE323] - C-terminal (ab32027) at 1/1000 dilution

Lane 1 : WEHI-231 (Mouse B cell lymphoma B lymphocyte) cell lysate

Lane 2 : F9 (Mouse embryonal carcinoma epithelial cell) cell lysate

Lane 3 : NIH/3T3 (Mouse embryonic fibroblast) cell lysate

Lane 4 : Mouse thymus lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution

Predicted band size: 99 kDa

Observed band size: 110 kDa

Blocking and diluting buffer and concentration: 5% NFDN/TBST

Exposure time: 30 seconds

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-CBL antibody [YE323] - C-terminal (ab32027)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors