abcam

Product datasheet

Anti-CD11c antibody [N418] ab33483

★★★★★ 14 Abreviews 77 References 2 图像

概述

产品名称 Anti-CD11c抗体[N418]

描述 亚美尼亚仓鼠单克隆抗体[N418] to CD11c

宿主Armenian hamster经测试应用适用于: Flow Cyt种属反应性与反应: Mouse

免疫原 Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.

阳性对照 IHC: mouse liver tissue, mouse spleen tissue Flow Cyt: Mouse splenocytes.

常规说明 Clone N418 has been reported to enhance antigen specific responses when used to target

dendritic cells in vivo.

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or

conjugation for your experiments, please contact orders@abcam.com.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

存储溶液 Preservative: 0.02% Sodium azide

Constituents: PBS, 6.97% L-Arginine

纯**度** Protein G purified

Primary antibody说明 Clone N418 has been reported to enhance antigen specific responses when used to target

dendritic cells in vivo.

克隆 单克隆

1

克隆编号 N418 骨髓瘤 Sp2/0 同种型 ΙgG

应用

靶标

Abpromise™承诺保证使用ab33483于以下的经测试应用 The Abpromise guarantee

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt	★★★★★ (1)	1/100. ab18479 - Armenian Hamster monoclonal lgG, is suitable for use as an isotype control with this antibody.

功能	Integrin alpha-X/beta-2 is a receptor for fibrinogen. It recognizes the sequence G-P-R in	
タJ RE	integrin alpha-7/beta-2 is a receptor for individuent. It recognizes the sequence G-F-1\forall in	

fibrinogen. It mediates cell-cell interaction during inflammatory responses. It is especially

important in monocyte adhesion and chemotaxis.

组织特异性 Predominantly expressed in monocytes and granulocytes.

序列相似性 Belongs to the integrin alpha chain family.


> Contains 7 FG-GAP repeats. Contains 1 VWFA domain.

结构域 The integrin I-domain (insert) is a VWFA domain. Integrins with I-domains do not undergo


protease cleavage.

细胞定位 Membrane.


图片



Flow Cytometry - Anti-CD11c antibody [N418] (ab33483)



Acquisition of >30,000 total events were collected. Gating strategy – events were collected with the forward and side light-scatter characteristics of viable lymphocytes.



Flow cytometry analysis of murine alveolar cells extracted using bronchoaveolar lavage in balb/c mice using 3mM PBS-EDTA, labelling CD11c with ab33483 at 1/2000, incubated for 20 mins at 20°C. Secondary used was **ab175680** at 1/2000. Gating strategy was against CD45+, C11b+, GR1-cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors