

Anti-CD200 / OX2 antibody [MRC OX90] ab33734

★★★★★ [1 Abreviews](#) [3 References](#) [1 图像](#)

概述

产品名称	Anti-CD200 / OX2抗体[MRC OX90]
描述	大鼠单克隆抗体[MRC OX90] to CD200 / OX2
宿主	Rat
经测试应用	适用于: IHC-FoFr
种属反应性	与反应: Mouse
免疫原	Fusion protein corresponding to Mouse CD200/ OX2. Database link: O54901
阳性对照	Flow Cyt: C57BL/6 splenocytes.
常规说明	<p>Spleen cells from immunised rats were fused with cells of the rat Y3 myeloma cell line.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	Preservative: 0.09% Sodium azide Constituent: PBS
纯度	Protein G purified
纯化说明	This purified antibody was prepared by affinity chromatography on Protein G from tissue culture supernatant.
Primary antibody说明	Spleen cells from immunised rats were fused with cells of the rat Y3 myeloma cell line.
克隆	单克隆

克隆编号	MRC OX90
同种型	IgG2a

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab33734于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-FoFr	★★★★★ (1)	Use at an assay dependent concentration.

靶标

功能	Costimulates T-cell proliferation. May regulate myeloid cell activity in a variety of tissues.
序列相似性	Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 Ig-like V-type (immunoglobulin-like) domain.
细胞定位	Cell membrane.

图片



Immunohistochemical analysis of murine brain tissue, staining CD200 / OX2 with ab33734.

Tissue was fixed with paraformaldehyde and blocked with 5% BSA + 5% serum for 1 hour at room temperature. Samples were incubated with primary antibody (1/600 in 5% BSA + 5% NGS) for 17 hours at 4°C. An AlexaFluor®555-conjugated goat anti-rat polyclonal IgG (1/200) was used as the secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors