abcam

Product datasheet

Anti-CD3 antibody [KT3] ab33429

★★★★★ 3 Abreviews 6 References 1 图像

概述

产品名称 Anti-CD3抗体[KT3]

描述 大鼠单克隆抗体[KT3] to CD3

宿主 Rat

经测试应用 适用于: IHC-Fr, ICC/IF, Flow Cyt

不适用于: IHC-P

种属反应性 与反应: Mouse

免疫原 Tissue, cells or virus corresponding to Mouse CD3. mature T cells (Mouse)

常规说明

ab33429 may be used to trigger proliferation and cytotoxicity of CD3 positive cells.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

存储溶液 pH: 7.40

Preservative: 0.09% Sodium azide

Constituent: PBS

纯**度** Protein G purified

纯**化**说明 Purified from tissue culture supernatant.

Primary antibody说明 ab33429 may be used to trigger proliferation and cytotoxicity of CD3 positive cells.

1

骨髓瘤 NS0

同种型 lgG2a

应用

The Abpromise guarantee Abpromise™承诺保证使用ab33429于以下的经测试应用


"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr	★★★★ (1)	Use at an assay dependent concentration.
ICC/IF	*****(1)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. <u>ab18450</u> - Rat monoclonal lgG2a, is suitable for use as an isotype control with this antibody.

应用说明

Is unsuitable for IHC-P.

图片



Immunohistochemistry (Frozen sections) - Anti-CD3 antibody [KT3] (ab33429)

Image courtesy of an anonymous Abreview

ab33429 staining CD3 in mouse spleen tissue by Immunohistochemistry (frozen sections). Tissue was fixed with acetone and then blocked with 5% serum for 1 hour at 25°C followed by incubation with the primary antibody at a 1/250 dilution for 12 hours at 4°C. A goat anti-rat IgG conjugated to Alexa Fluor®488 was used as secondary antibody at a 1/250 dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors