


### Anti-CD34 antibody [EP373Y] ab81289

重组 RabMAb

★★★★★ **55 Abreviews** **499 References** **13 图像**

#### 概述

产品名称	Anti-CD34抗体[EP373Y]
描述	兔单克隆抗体[EP373Y] to CD34
宿主	Rabbit
经测试应用	<b>适用于:</b> IHC-P, ICC/IF, IP, IHC-Fr, Flow Cyt (Intra) <b>不适用于:</b> WB
种属反应性	<b>与反应:</b> Mouse, Rat, Human <b>预测可用于:</b> Sheep, Dog, Pig, African bush elephant 
免疫原	Synthetic peptide within Human CD34 aa 350 to the C-terminus (C terminal). The exact sequence is proprietary. Database link: <b>P28906</b>
阳性对照	Flow Cyt (intra): TF-1 cells. IHC-P: Human kidney tissue; Mouse normal brain, prostate and kidney tissues; Rat kidney tissue. ICC/IF: HUVEC cells; Human embryonic stem cell-derived endothelial cells. IHC-Fr: Mouse and rat lung tissue. Rat kidney. IP: TF-1 cell lysate.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <b><a href="#">see here</a></b> . Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <b><a href="#">RabMAb<sup>®</sup> patents</a></b> .

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 1.15 x 10 <sup>-10</sup> M





[Learn more about K<sub>D</sub>](#)

存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EP373Y
同种型	IgG

## 应用

**The Abpromise guarantee** **Abpromise™** 承诺保证使用 ab81289 于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度; 实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
IHC-P	★★★★★ (32)	1/2500 - 1/5000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. See <a href="#">IHC antigen retrieval protocols</a> .
ICC/IF	★★★★★ (5)	1/100 - 1/250.
IP		1/30.
IHC-Fr	★★★★★ (7)	1/50.
Flow Cyt (Intra)		1/50.


应用说明 Is unsuitable for WB.

## 靶标


功能	Possible adhesion molecule with a role in early hematopoiesis by mediating the attachment of stem cells to the bone marrow extracellular matrix or directly to stromal cells. Could act as a scaffold for the attachment of lineage specific glycans, allowing stem cells to bind to lectins expressed by stromal cells or other marrow components. Presents carbohydrate ligands to selectins.
组织特异性	Selectively expressed on hematopoietic progenitor cells and the small vessel endothelium of a variety of tissues.
序列相似性	Belongs to the CD34 family.
发展阶段	On early hematopoietic progenitor cells.
翻译后修饰	Highly glycosylated. Phosphorylated on serine residues by PKC.
细胞定位	Membrane.



Intracellular Flow Cytometry analysis of TF-1 (human erythroleukemia) cells labeling CD34 with purified ab81289 at 1/50 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling CD34 with purified ab81289 at 1/2500. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) was used as the secondary antibody at 1/500. Negative control using PBS instead of primary antibody. Counter stained with Hematoxylin.



Immunocytochemistry/ Immunofluorescence - Anti-CD34 antibody [EP373Y] (ab81289)


Immunocytochemistry/Immunofluorescence analysis of HUVEC (Human umbilical vein endothelial cell line) cells labelling CD34 with purified ab81289 at 1/100. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG (**ab150077**) (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counter stain.

Control: Secondary antibody only, **ab150120**, an Alexa Fluor® 488-conjugated goat anti-mouse IgG (1/500).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD34 antibody [EP373Y] (ab81289)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat kidney tissue labelling CD34 with purified ab81289 at 1/2500. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) was used as the secondary antibody at 1/500. Negative control using PBS instead of primary antibody. Counter stained with Hematoxylin.




Immunohistochemistry (Frozen sections) - Anti-CD34 antibody [EP373Y] (ab81289)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat lung tissue labeling CD34 with ab81289 at 1/50 (11.04 µg/mL) dilution followed by **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on rat lung. is observed. The nuclear counterstain was DAPI (Blue).

Secondary antibody control: Secondary antibody is **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Frozen sections) - Anti-CD34 antibody [EP373Y] (ab81289)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen Rat kidney tissue labeling CD34 with ab81289 at 1/50 (11.04 µg/mL) dilution followed by **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution (Green). Positive staining on rat kidney. is observed. The nuclear counterstain was DAPI (Blue).

Secondary antibody control: Secondary antibody is **ab150077** Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1000 dilution.


Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD34 antibody [EP373Y] (ab81289)

This image is courtesy of an Abreview submitted by Mr. Rudolf Jung.

Paraformaldehyde-fixed, paraffin-embedded mouse kidney tissue stained for CD34 using ab81289 at 1/200 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD34 antibody [EP373Y] (ab81289)

IHC image of CD34 staining in Mouse normal brain formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with unpurified ab81289, 1/250 dilution, for 15 mins at room temperature. A Goat anti-Rabbit biotinylated secondary antibody was used to detect the primary, and visualized using an HRP conjugated ABC system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.





Immunoprecipitation - Anti-CD34 antibody [EP373Y]  
(ab81289)


CD34 was immunoprecipitated from TF-1 (Human bone marrow erythroleukemia cells) cells using purified ab81289 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab81289. Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated was used as secondary antibody at 1/1000 dilution. Blocking and dilution buffer and concentration: 5% NFDM/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD34 antibody [EP373Y]  
(ab81289)

This image is courtesy of an anonymous Abreview.

Paraformaldehyde-fixed, paraffin-embedded mouse prostate tissue stained for CD34 using ab81289 at 1/150 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence - Anti-CD34 antibody [EP373Y] (ab81289)

Image from Földes Get al, PLoS One. 2010 May 5;5(5):e10501, Fig 3.

Immunocytochemistry/Immunofluorescence analysis of human embryonic stem cell-derived endothelial cells labeling CD34 with unpurified ab81289 at 1/200. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.2% aTriton X-100. An Alexa Fluor® 647-conjugated secondary antibody was used at a 1/400 dilution. DAPI (blue) was used as the nuclear counter stain.

OI-RD Scanning - Anti-CD34 antibody [EP373Y]  
(ab81289)

Equilibrium disassociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)

Why choose a  
recombinant antibody?



**Research with  
confidence**  
Consistent and  
reproducible results



**Long-term and  
scalable supply**  
Recombinant  
technology



**Success from the  
first experiment**  
Confirmed  
specificity



**Ethical standards  
compliant**  
Animal-free  
production

Anti-CD34 antibody [EP373Y] (ab81289)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors