

Anti-CD5 antibody [EP2952] ab75877

重组 RabMAb


2 References **9 图像**

概述

产品名称	Anti-CD5抗体[EP2952]
描述	兔单克隆抗体[EP2952] to CD5
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P 不适用于: Flow Cyt
种属反应性	与反应: Human
免疫原	Synthetic peptide within Human CD5 aa 50-150. The exact sequence is proprietary.
阳性对照	WB: Human lymph node lysate IHC-P: Human tonsil tissue. IP: Human lymph node lysate
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
解离常数 (K_D)	K _D = 1.70 x 10 ⁻¹¹ M



[Learn more about K_D](#)

存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.5% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EP2952
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab75877 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB		1/10000 - 1/50000. Predicted molecular weight: 55 kDa.
IP		1/50.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

应用说明 Is unsuitable for Flow Cyt.

靶标

功能	May act as a receptor in regulating T-cell proliferation. CD5 interacts with CD72/LYB-2.
序列相似性	Contains 3 SRCR domains.
细胞定位	Membrane.

图片



Immunoprecipitation - Anti-CD5 antibody [EP2952]
(ab75877)


CD5 was immunoprecipitated from 0.35 mg Human lymph node lysate 10 μ g with 75877 at 1/20 dilution (0.6 μ g) . VeriBlot for IP Detection Reagent (HRP)(**ab131366**) was used at 1/5000 dilution.

Lane 1: Human lymph node lysate 10 μ g

Lane 2: ab75877 IP in Human lymph node lysate

Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab75877 in Human lymph node lysate

Blocking and dilution buffer and concentration: 5% NFDN/TBST.



Western blot - Anti-CD5 antibody [EP2952]
(ab75877)


Anti-CD5 antibody [EP2952] (ab75877) at 1/1000 dilution (Purified)
+ Human lymph node lysate at 15 μ g

Secondary

Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 55 kDa


Observed band size: 67 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD5 antibody [EP2952]
(ab75877)

ab75877 showing positive staining in Normal colon lymphocytes tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD5 antibody [EP2952] (ab75877)

ab75877 showing negative staining in Normal brain tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD5 antibody [EP2952] (ab75877)

ab75877 showing negative staining in Normal kidney tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD5 antibody [EP2952] (ab75877)

ab75877 showing negative staining in Skeletal muscle tissue.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD5 antibody [EP2952] (ab75877)

ab75877 showing positive staining in Normal spleen tissue.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Ox-LD Scanning - Anti-CD5 antibody [EP2952] (ab75877)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?

- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-CD5 antibody [EP2952] (ab75877)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors