

Anti-CD8 alpha antibody [14] ab20133

2 References

概述

产品名称	Anti-CD8 alpha抗体[14]
描述	小鼠单克隆抗体[14] to CD8 alpha
宿主	Mouse
特异性	Specific to CD8 alpha, ab20133 can be used for the detection of immunoregulatory T cell subset imbalances in autoimmune disorders and immunodeficiency states, categorisation of T cell acute lymphoblastic leukaemia and lymphoblastic lymphoma in conjunction with other antigens and determination of T4/T8 ratios in immune deficiencies, rheumatoid arthritis, multiple sclerosis, etc.
经测试应用	适用于: ELISA, Flow Cyt, IHC, IP
种属反应性	与反应: Human
免疫原	Tissue, cells or virus corresponding to CD8 alpha. Fibronectin purified monocytes.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium azide Constituent: 99.98% PBS
纯度	Protein A/G purified
克隆	单克隆
克隆编号	14
骨髓瘤	unknown
同种型	IgG1

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab20133于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. FACS
IHC		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

靶标

功能	Identifies cytotoxic/suppressor T-cells that interact with MHC class I bearing targets. CD8 is thought to play a role in the process of T-cell mediated killing. CD8 alpha chains binds to class I MHC molecules alpha-3 domains.
疾病相关	Defects in CD8A are a cause of familial CD8 deficiency (CD8 deficiency) [MIM:608957]. Familial CD8 deficiency is a novel autosomal recessive immunologic defect characterized by absence of CD8+ cells, leading to recurrent bacterial infections.
序列相似性	Contains 1 Ig-like V-type (immunoglobulin-like) domain.
翻译后修饰	All of the five most carboxyl-terminal cysteines form inter-chain disulfide bonds in dimers and higher multimers, while the four N-terminal cysteines do not.
细胞定位	Secreted and Cell membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors