

Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free ab213216

敲除验证
重组
RabMAb

3 References 10 图像

概述


产品名称	Anti-Cdk4抗体[EPR4513-32-7] - BSA and Azide free
描述	兔单克隆抗体[EPR4513-32-7] to Cdk4 - BSA and Azide free
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, ICC/IF, IHC-P
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HAP1 and HeLa cell lysates, MCF7 membrane extract lysate (ab29539). Flow Cyt (intra): MCF-7 IHC-P: Human cervix carcinoma and human urothelial carcinoma. ICC/IF: MCF-7 and HAP1.
常规说明	<p>ab213216 is the carrier-free version of ab108357.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
解离常数 (K_D)	$K_D = 1.86 \times 10^{-11}$ M



存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR4513-32-7
同种型	IgG

应用

The Abpromise guarantee [Abpromise™](#) 承诺保证使用 ab213216 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Detects a band of approximately 34 kDa (predicted molecular weight: 34 kDa).
ICC/IF		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols .

靶标

功能	Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit
----	---

members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex.

疾病相关

Defects in CDK4 are a cause of susceptibility to cutaneous malignant melanoma type 3 (CMM3) [MIM:609048]. Malignant melanoma is a malignant neoplasm of melanocytes, arising de novo or from a pre-existing benign nevus, which occurs most often in the skin but also may involve other sites.

序列相似性

Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily.

Contains 1 protein kinase domain.


翻译后修饰

Phosphorylation at Thr-172 is required for enzymatic activity. Phosphorylated, in vitro, at this site by CCNH-CDK7, but, in vivo, appears to be phosphorylated by a proline-directed kinase. In the cyclin D-CDK4-CDKN1B complex, this phosphorylation and consequent CDK4 enzyme activity, is dependent on the tyrosine phosphorylation state of CDKN1B. Thus, in proliferating cells, CDK4 within the complex is phosphorylated on Thr-172 in the T-loop. In resting cells, phosphorylation on Thr-172 is prevented by the non-tyrosine-phosphorylated form of CDKN1B.

细胞定位

Cytoplasm. Nucleus. Membrane. Cytoplasmic when non-complexed. Forms a cyclin D-CDK4 complex in the cytoplasm as cells progress through G(1) phase. The complex accumulates on the nuclear membrane and enters the nucleus on transition from G(1) to S phase. Also present in nucleoli and heterochromatin lumps. Colocalizes with RB1 after release into the nucleus.

图片



Western blot - Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

All lanes : Anti-Cdk4 antibody [EPR4513-32-7] ([ab108357](#)) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CDK4 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 34 kDa

Observed band size: 34 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab108357](#)).

Lanes 1- 2: Merged signal (red and green). Green - **ab108357** observed at 34 kDa. Red - Anti-Vinculin antibody [VIN-54] observed at 124 kDa.

ab108357 was shown to react with Cdk4 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line **ab255378** (knockout cell lysate **ab263780**) was used. Wild-type HeLa and CDK4 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. **ab108357** and Anti-Vinculin antibody [VIN-54] overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Flow Cytometry (Intracellular) - Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

ab108357 staining CDK4 in the human cell line MCF-7 (human breast carcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde, permeabilized with 90% methanol and the sample was incubated with the primary antibody at a dilution of 1/20. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isoytype control: Rabbit monoclonal IgG (Black)

Unlabeled control: Cell without incubation with primary antibody and secondary antibody (Blue)


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk4 antibody
[EPR4513-32-7] - BSA and Azide free (ab213216)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervix carcinoma tissue labeling Cdk4 with purified **ab108357** at 1/100. Heat mediated antigen retrieval was performed using Tris/EDTA buffer (pH 9). **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L), was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).




Immunocytochemistry/ Immunofluorescence - Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

Immunocytochemical analysis of MCF7 cells, labeling Cdk4 with purified **ab108357** at 1/100. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500), was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.

Control 1: primary antibody (1/100) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).



Lanes 1-2 : Anti-Cdk4 antibody [EPR4513-32-7] ([ab108357](#)) at 1/1000 dilution

Lanes 3-4 : Anti-beta Actin antibody [mAbcam 8226] - Loading Control ([ab8226](#)) at 1/1000 dilution

Lanes 1 & 3 & 5 : Wild-type HAP1 cell lysate

Lanes 2 & 4 & 6 : CDK4 knockout HAP1 cell lysate


Lysates/proteins at 20 µg per lane.

Predicted band size: 34 kDa

This data was developed using the same antibody clone in a different buffer formulation ([ab108357](#)).

Lanes 5 and 6: Merged signal (red and green). Green - [ab108357](#) observed at 34kDa. Red - loading control to beta actin observed at 40kDa.


[ab108357](#) was shown to specifically react with CDK4 in wild-type HAP1 cells. No band was observed when CDK4 knockout samples were examined. Wild-type and CDK4 knockout samples were subjected to SDS-PAGE. [ab108357](#) and [ab8226](#) (loading control to beta actin) were both diluted at 1/1000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-Cdk4 antibody
[EPR4513-32-7] - BSA and Azide free (ab213216)

Intracellular Flow Cytometry analysis of MCF7 cells labelling Cdk4 with purified **ab108357** at 1/40 (red). Cells were fixed with 2% paraformaldehyde. A FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabelled control, cells without incubation with primary and secondary antibodies.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).



Immunocytochemistry/ Immunofluorescence - Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

ab108357 staining Cdk4 in wild-type HAP1 cells (top panel) and Cdk4 knockout HAP1 cells (bottom panel). The cells were fixed with 4% formaldehyde (10min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with **ab108357** at 1/500 dilution and **ab195889** at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green). Nuclear DNA was labelled in blue with DAPI.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk4 antibody
[EPR4513-32-7] - BSA and Azide free (ab213216)

Immunohistochemical analysis of formalin/PFA-fixed paraffin-embedded human urothelial carcinoma tissue labelling Cdk4 with unpurified **ab108357** at a dilution of 1/100.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab108357**).



SPR Scanning - Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab108357](#)).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Cdk4 antibody [EPR4513-32-7] - BSA and Azide free (ab213216)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors