

Anti-Cdk9 antibody [EPR3119Y] ab76320

重组 RabMAb

★★★★☆ **3 Abreviews** **14 References** **15 图像**

概述

产品名称	Anti-Cdk9抗体[EPR3119Y]
描述	兔单克隆抗体[EPR3119Y] to Cdk9
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P, ICC/IF, Flow Cyt (Intra)
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HT-29, NIH/3T3, C6, HL-60, HeLa or 293 cell lysates. IHC-P: Human breast carcinoma, Human Bladder, mouse testis tissue. ICC/IF: HT-29 cells Flow Cyt (intra): HeLa cells IP: HeLa cells
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR3119Y
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab76320于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/5000. Predicted molecular weight: 43 kDa. For unpurified use as 1/1000 - 1/2000
IP		1/20. For unpurified use as 1/40
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .
ICC/IF	★★★★★ (1)	1/50. For unpurified use as 1/100 - 1/250
Flow Cyt (Intra)		1/60.

靶标

功能	Member of the cyclin-dependent kinase pair (CDK9/cyclin-T) complex, also called positive transcription elongation factor b (P-TEFb), which facilitates the transition from abortive to production elongation by phosphorylating the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP II), SUPT5H and RDBP. The CDK9/cyclin-K complex has also a kinase activity toward CTD of RNAP II and can substitute for P-TEFb in vitro.
组织特异性	Ubiquitous.
序列相似性	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily. Contains 1 protein kinase domain.
细胞定位	Nucleus.

图片



Western blot - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

All lanes : Anti-Cdk9 antibody [EPR3119Y] (ab76320) at 1/5000 dilution (Purified)

Lane 1 : HT-29 (Human colorectal adenocarcinoma epithelial cell) whole cell lysates

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates

Lane 3 : C6 (Rat glial tumor glial cell) whole cell lysates


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 43 kDa

Observed band size: 43 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat testis tissue sections labeling Cdk9 with Purified ab76320 at 1:100 dilution (1.08 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunocytochemistry/ Immunofluorescence - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Immunocytochemistry/ Immunofluorescence analysis of HT-29 (Human colorectal adenocarcinoma epithelial cell) cells labeling Cdk9 with Purified ab76320 at 1:50 dilution (2.2 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse testis tissue sections labeling Cdk9 with Purified ab76320 at 1:100 dilution (1.08 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human bladder carcinoma tissue sections labeling Cdk9 with Purified ab76320 at 1:100 dilution (1.08 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Flow Cytometry (Intracellular) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Cdk9 with Purified ab76320 at 1/60 dilution (10 µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunoprecipitation - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

ab76320 (purified) at 1:20 dilution (0.5µg) immunoprecipitating Cdk9 in HeLa whole cell lysate.
 Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg
 Lane 2 (+): ab76320 & HeLa whole cell lysate
 Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab76320 in HeLa whole cell lysate
 For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.
 Blocking and diluting buffer: 5% NFD/MTBST.



Western blot - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

All lanes : Anti-Cdk9 antibody [EPR3119Y] (ab76320) at 1/2000 dilution (Unpurified)

Lane 1 : HT-29 cell lysate

Lane 2 : HL-60 cell lysate

Lane 3 : HeLa cell lysate


Lane 4 : 293 cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat anti-rabbit HRP at 1/1000 dilution


Predicted band size: 43 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Image courtesy of an Abreview submitted by Dr. Kirk McManus, Univ. of Manitoba/Cancer Care MCB, Canada


Unpurified ab76320 (1/200) staining Cdk9 in HeLa cells (green). Cells were fixed in paraformaldehyde, permeabilised in 0.5% Triton X100/ PBS and counterstained with DAPI in order to highlight the nucleus (red). For further experimental details please refer to Abreview.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma using unpurified ab76320 at 1/100 dilution.


Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Unpurified ab76320 showing positive staining in Cervical carcinoma tissue.


Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Unpurified ab76320 showing positive staining in Colonic adenocarcinoma tissue.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Unpurified ab76320 showing positive staining in Lung adenocarcinoma tissue.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Unpurified ab76320 showing positive staining in Normal tonsil tissue.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Cdk9 antibody [EPR3119Y] (ab76320)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors