


### Anti-Cip4 antibody [EPR1965] ab108313

敲除验证
重组
RabMAb

[1 References](#)
[4 图像](#)

#### 概述

产品名称	Anti-Cip4抗体[EPR1965]
描述	兔单克隆抗体[EPR1965] to Cip4
宿主	Rabbit
特异性	Recent WB re-tests performed by our lab suggest to use a higher dilution for mouse samples. The tested dilutions were 1:500 and 1:2000, where the signal detected was very strong. Human and rat samples worked fine in these dilutions.
经测试应用	<b>适用于:</b> WB, IHC-P <b>不适用于:</b> Flow Cyt, ICC/IF or IP
种属反应性	<b>与反应:</b> Human <b>预测可用于:</b> Mouse, Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HEK293T, JAR, HepG2, TF1, and HeLa cell lysates. IHC-P: Human stomach tissue.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.20 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
纯度	Tissue culture supernatant

克隆	单克隆
克隆编号	EPR1965
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise™承诺保证使用ab108313于以下的经测试应用**

**“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。**

应用	Ab评论	说明
<b>WB</b>		1/1000 - 1/10000. Predicted molecular weight: 68 kDa.
<b>IHC-P</b>		1/100 - 1/250. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.

**应用说明** Is unsuitable for Flow Cyt, ICC/IF or IP.

## 靶标

**功能** Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.


组织特异性	Expressed in brain, colon, heart, kidney, liver, lung, megakaryocyte, ovary, pancreas, peripheral blood lymphocytes, placenta, prostate, skeletal muscle, small intestine, spleen, testis, thymus and trachea.
-------	--

序列相似性	<p>Belongs to the FNBP1 family.</p> <p>Contains 1 FCH domain.</p> <p>Contains 1 REM (Hr1) repeat.</p> <p>Contains 1 SH3 domain.</p>
-------	---

**翻译后修饰** Tyrosine phosphorylated. Also phosphorylated by PKA.

**细胞定位** Cytoplasm > perinuclear region and Cytoplasm > cytoskeleton. Cytoplasm > cell cortex. Lysosome. Golgi apparatus. Cell membrane. Cell projection > phagocytic cup. Translocates to the plasma membrane in response to insulin stimulation, and this may require active RHOQ (By similarity). Localizes to cortical regions coincident with F-actin, to lysosomes and to sites of phagocytosis in macrophages. Also localizes to the Golgi, and this requires AKAP9.

图片



Western blot - Anti-Cip4 antibody [EPR1965] (ab108313)

**All lanes :** Anti-Cip4 antibody [EPR1965] (ab108313) at 1/1000 dilution

**Lane 1 :** Wild-type HEK293T cell lysate

**Lane 2 :** TRIP10 knockout HEK293T cell lysate

**Lane 3 :** JAR cell lysate

**Lane 4 :** HepG2 cell lysate

Lysates/proteins at 20 µg per lane.

## Secondary


**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 68 kDa

**Observed band size:** 75 kDa

**Lanes 1-4:** Merged signal (red and green). Green - ab108313 observed at 75 kDa. Red - loading control [ab8245](#) observed at 36 kDa.


ab108313 Anti-Cip4 antibody [EPR1965] was shown to specifically react with Cip4 in wild-type HEK293T cells. Loss of signal was observed when knockout cell line [ab266428](#) (knockout cell lysate [ab258251](#)) was used. Wild-type and Cip4 knockout samples were subjected to SDS-PAGE. ab108313 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cip4 antibody [EPR1965] (ab108313)

ab108313 at 1/100 dilution staining Cip4 in paraffin embedded Human stomach tissue.

Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol.



Western blot - Anti-Cip4 antibody [EPR1965] (ab108313)

**All lanes :** Anti-Cip4 antibody [EPR1965] (ab108313) at 1/1000 dilution

**Lane 1 :** JAR cell lysate

**Lane 2 :** HepG2 cell lysate

**Lane 3 :** TF1 cell lysate

**Lane 4 :** HeLa cell lysate

Lysates/proteins at 10 µg per lane.

**Predicted band size:** 68 kDa

Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Cip4 antibody [EPR1965] (ab108313)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors