

Anti-Cox-3 antibody ab37269

1 References [1 图像](#)

概述

产品名称	Anti-Cox-3抗体
描述	兔多克隆抗体to Cox-3
宿主	Rabbit
经测试应用	适用于: WB
种属反应性	与反应: Mouse
免疫原	Synthetic peptide corresponding to residues from the N-terminus of Cox-3 (Mouse).
阳性对照	mouse small intestinal tissue
常规说明	<p>This product is manufactured by BioVision, an Abcam company and was previously called 3687 Cox-3 Antibody. 3687-100 is the same size as the 100 µg size of ab37269.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.02% Thimerosal (merthiolate) Constituents: 1% BSA, 50% Glycerol (glycerin, glycerine), PBS
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用 ab37269 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
WB		Use at an assay dependent concentration. Predicted molecular weight: 69 kDa.

靶标


相关性

COX proteins are membrane-associated heme proteins that have cyclooxygenase and peroxidase activities. These enzymes are targets of NSAID (nonsteroidal anti-inflammatory drugs) such as aspirin. Prostaglandins (PGs) formed by the enzymatic activity of COX-1 are primarily involved in the regulation of homeostatic functions throughout the body, whereas PGs formed by COX-2 primarily mediate pain, fever, and inflammation. COX-1 is constitutively expressed, with particularly high expression in gastrointestinal tissues. COX-2 is induced by cytokines and mitogens and is likely to play a role in inflammatory diseases such as rheumatoid arthritis. In rodents and humans, COX-3 encodes proteins with completely different amino acid sequences from COX-1 or COX-2, and without COX activity. COX-3 is thought to be a splice variant of COX-1 which retains intron one. It has been suggested that COX-3 may be the key to unlocking the mechanism of action of acetaminophen.

细胞定位

Membrane-associated; Microsomal membrane.

图片



Anti-Cox-3 antibody (ab37269) (0.5–4 µg/ml) + Mouse small intestine tissue lysate

Predicted band size: 69 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors