

Anti-Cyclin A2 antibody [Y193] ab32386

重组 RabMAb

★★★★★ **7 Abreviews** **26 References** **7 图像**

概述

产品名称	Anti-Cyclin A2抗体[Y193]
描述	兔单克隆抗体[Y193] to Cyclin A2
宿主	Rabbit
经测试应用	适用于: IHC-P, IP, WB, ICC/IF
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa, HepG2, HEK293, NIH/3T3 cell lysates. IHC-P: Skin and colon carcinoma. ICC/IF: MCF7 cells. IP: HeLa cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 49% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆

克隆编号 Y193

同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab32386于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (1)	1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
IP		1/40 - 1/50.
WB	★★★★★ (2)	1/10000. Detects a band of approximately 55 kDa (predicted molecular weight: 48 kDa). For unpurified, use 1/1000.
ICC/IF	★★★★★ (3)	1/100 - 1/250.

靶标


功能 Essential for the control of the cell cycle at the G1/S (start) and the G2/M (mitosis) transitions.

序列相似性 Belongs to the cyclin family. Cyclin AB subfamily.

发展阶段 Accumulates steadily during G2 and is abruptly destroyed at mitosis.

细胞定位 Nucleus. Cytoplasm. Cytoplasmic when associated with SCAPER.

图片




Anti-Cyclin A2 antibody [Y193] (ab32386) at 1/10000 dilution (purified) + NIH/3T3 cell lysate at 20 µg

Secondary
Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 48 kDa
Observed band size: 55 kDa

Blocking buffer: 5% NFDM/TBST
Dilution buffer: 5% NFDM/TBST



All lanes : Anti-Cyclin A2 antibody [Y193] (ab32386) at 1/10000 dilution (purified)

Lane 1 : HepG2 cell lysate

Lane 2 : HEK293 cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 48 kDa


Observed band size: 55 kDa

Blocking buffer: 5% NFDM/TBST

Dilution buffer: 5% NFDM/TBST




ab32386 (purified) at 1/40 immunoprecipitating Cyclin A2 in 10 µg HeLa cell lysate (Lanes 1 and 2, observed at 55 kDa). Lane 3 - Rabbit monoclonal IgG ([ab172730](#)). For western blotting, HRP Veriblot for IP Detection Reagent ([ab131366](#)) was used for detection (1/10 000). Blocking buffer and concentration: 5% NFDM/TBST Dilution buffer and concentration: 5% NFDM/TBST




Immunocytochemistry/ Immunofluorescence - Anti-Cyclin A2 antibody [Y193] (ab32386)

Immunofluorescence staining of MCF7 cells with purified ab32386 at a working dilution of 1/100, counter-stained with DAPI. The secondary antibody was Alexa Fluor® 488 goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. **ab7291**, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with **ab150120** (Alexa Fluor® 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4 % PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab32386 was used at a dilution of 1/500 followed by an Alexa Fluor® 594 goat anti-mouse antibody (**ab150120**) at a dilution of 1/500. For negative control 2, **ab7291** (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor® 488 goat anti-rabbit antibody (**ab150077**) at a dilution of 1/400.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cyclin A2 antibody [Y193] (ab32386)

Immunohistochemical staining of paraffin embedded human colon carcinoma with purified ab32386 at a working dilution of 1/500. The secondary antibody used is **ab97051**, a goat anti-rabbit IgG (H&L) at a dilution of 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.







Anti-Cyclin A2 antibody [Y193] (ab32386) at 1/1000 dilution
(unpurified) + HeLa cell lysate

Predicted band size: 48 kDa

Observed band size: 55 kDa

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-Cyclin A2 antibody [Y193] (ab32386)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors