abcam

Product datasheet

Anti-CYP27A1 antibody [EPR7529] ab126785



重组 RabMAb

★★★★★ 6 Abreviews 31 References 7 图像

概述

产品名称 Anti-CYP27A1抗体[EPR7529]

描述 兔单克隆抗体[EPR7529] to CYP27A1

宿主 Rabbit

经测试应用 适用于: Flow Cyt (Intra), WB, IHC-P, ICC/IF

不适用干: IP

种属反应性 与反应: Mouse. Human

预测可用于: Rat 📤

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

阳性对照 WB: HepG2, Caco 2, human fetal liver and mouse liver lysates. IHC-P: Human kidney and liver

tissues. ICC/IF: HepG2 cells. Flow Cyt (intra): HepG2 cells.

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.


Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® patents.

性能

形式

存放说明 Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.

 $K_D = 6.00 \times 10^{-11} M$ 解离常数(Kn)



Learn more about KD

存储溶液 pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture

supernatant

纯**度** Protein A purified

克隆 单克隆

克隆编号 EPR7529

同种型 IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab126785于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|------------------|------------------|---|
| Flow Cyt (Intra) | | 1/120. |
| WB | ★★★★★ (3) | 1/1000 - 1/10000. Predicted molecular weight: 60 kDa. |
| IHC-P | ★★★★ (2) | 1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |
| ICC/IF | | 1/50 - 1/100. |

应用说明 Is unsuitable for IP.

靶标

功能 Catalyzes the first step in the oxidation of the side chain of sterol intermediates; the 27-

hydroxylation of 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol. Has also a vitamin D3-25-

hydroxylase activity.


通路 Hormone biosynthesis; cholecalciferol biosynthesis.

疾病相关 Cerebrotendinous xanthomatosis

序列相似性 Belongs to the cytochrome P450 family.

细胞定位 Mitochondrion membrane.

图片



Western blot - Anti-CYP27A1 antibody [EPR7529] (ab126785)

All lanes : Anti-CYP27A1 antibody [EPR7529] (ab126785) at 1/1000 dilution

Lane 1 : HepG2 lysate Lane 2 : Caco 2 lysate

Lane 3: Human fetal liver lysate


Lane 4: Mouse liver lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes: HRP labelled goat anti-rabbit at 1/2000 dilution


Predicted band size: 60 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CYP27A1 antibody
[EPR7529] (ab126785)


ab126785, at a 1/100 dilution, staining CYP27A1 in paraffin embedded Human kidney tissue by Immunohistochemistry.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-CYP27A1 antibody [EPR7529] (ab126785)

ab126785, at a 1/50 dilution, staining CYP27A1 in HepG2 cells by Immunofluorescence.



Flow Cytometry (Intracellular) - Anti-CYP27A1 antibody [EPR7529] (ab126785)


Intracellular Flow Cytometry analysis of HepG2 (human hepatocellular carcinoma) cells labeling CYP27A1 (red) with ab126785 at a 1/120 dilution. Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti-rabbit lgG (Alexa Fluorr® 488) (ab150077) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal lgG (ab172730). Blue (unlabeled control) - Cells without incubation with primary and secondary antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-CYP27A1 antibody
[EPR7529] (ab126785)

ab126785, at a 1/100 dilution, staining CYP27A1 in paraffin embedded Human liver tissue by Immunohistochemistry.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Ol-RD Scanning - Anti-CYP27A1 antibody [EPR7529] (ab126785)

Equilibrium disassociation constant (K_D) Learn more about K_D

Click here to learn more about KD



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors