

# Anti-Cytochrome C antibody [EPR1327] - BSA and Azide free ab218312

 **RabMAb**


**8 References**   **9 图像**

### 概述

产品名称	Anti-Cytochrome C抗体[EPR1327] - BSA and Azide free
描述	兔单克隆抗体[EPR1327] to Cytochrome C - BSA and Azide free
宿主	Rabbit
经测试应用	<b>适用于:</b> WB, IP, IHC-P, ICC/IF
种属反应性	<b>与反应:</b> Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Molt4, SH-SY5Y, Human heart, Human kidney and Human spleen lysates. Human kidney tissue.
常规说明	<p>ab218312 is the carrier-free version of <a href="#">ab133504</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <b>conjugation kits</b> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 1.29 x 10 <sup>-10</sup> M



存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR1327
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™** 承诺保证使用 ab218312 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB		Use at an assay dependent concentration. Detects a band of approximately 14 kDa (predicted molecular weight: 11 kDa).
IP		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <a href="#">IHC antigen retrieval protocols</a> .
ICC/IF		Use at an assay dependent concentration.

靶标

功能	<p>Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.</p> <p>Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.</p>
----	--


疾病相关	Defects in CYCS are the cause of thrombocytopenia type 4 (THC4) [MIM:612004]; also known as autosomal dominant thrombocytopenia type 4. Thrombocytopenia is the presence of relatively few platelets in blood. THC4 is a non-syndromic form of thrombocytopenia. Clinical manifestations of thrombocytopenia are absent or mild. THC4 may be caused by dysregulated platelet formation.
序列相似性	Belongs to the cytochrome c family.
翻译后修饰	Binds 1 heme group per subunit.
细胞定位	Mitochondrion matrix.

图片




Immunohistochemical staining of paraffin embedded human cervical carcinoma with purified **ab133504** at a working dilution of 1 in 500. The secondary antibody used is a HRP goat anti-rabbit H+L (**ab97051**). The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).



Immunohistochemical staining of paraffin embedded mouse liver with purified **ab133504** at a working dilution of 1 in 500. The secondary antibody used is a HRP goat anti-rabbit H+L (**ab97051**). The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytochrome C antibody [EPR1327] - BSA and Azide free (ab218312)

Immunohistochemical staining of paraffin embedded rat kidney with purified **ab133504** at a working dilution of 1 in 500. The secondary antibody used is a HRP goat anti-rabbit H+L (**ab97051**). The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).



Immunocytochemistry/ Immunofluorescence - Anti-Cytochrome C antibody [EPR1327] - BSA and Azide free (ab218312)

Immunofluorescence staining of SH-SY5Y cells with purified **ab133504** at a working dilution of 1 in 100, counter-stained with DAPI. The secondary antibody was Alexa Fluor<sup>®</sup> 488 goat anti rabbit (**ab150077**), used at a dilution of 1 in 500. **ab7291** was used to stain tubulin, and this is shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative control is shown in bottom middle and right hand panels - for the negative controls, purified **ab133504** was used at a dilution of 1/200 followed by an Alexa Fluor<sup>®</sup> 594 goat anti-mouse antibody at a dilution of 1/500.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).




Immunoprecipitation - Anti-Cytochrome C antibody  
[EPR1327] - BSA and Azide free (ab218312)

**ab133504** (purified) at 1/30 immunoprecipitating Cytochrome C in Molt-4 cells (Lane 1). For western blotting, a HRP-conjugated goat anti-rabbit (H+L), was used as the secondary antibody (1/1000).

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).




OxLD Scanning - Anti-Cytochrome C antibody  
[EPR1327] - BSA and Azide free (ab218312)

Equilibrium disassociation constant ( $K_D$ )

Learn more about  $K_D$

[Click here to learn more about  \$K\_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytochrome C antibody  
[EPR1327] - BSA and Azide free (ab218312)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labelling Cytochrome C with unpurified **ab133504** at 1/250 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.





Immunofluorescent analysis of HeLa cells labelling Cytochrome C with unpurified **ab133504** at 1/100 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab133504**).

Immunocytochemistry/ Immunofluorescence - Anti-Cytochrome C antibody [EPR1327] - BSA and Azide free (ab218312)

Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-Cytochrome C antibody [EPR1327] - BSA and Azide free (ab218312)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors