

Anti-Cytokeratin 17 antibody - Cytoskeleton Marker ab53707

★★★★★ [2 Abreviews](#) [36 References](#) [3 图像](#)

概述

| | |
|-------|---|
| 产品名称 | Anti-Cytokeratin 17抗体- Cytoskeleton Marker |
| 描述 | 兔多克隆抗体to Cytokeratin 17 - Cytoskeleton Marker |
| 宿主 | Rabbit |
| 经测试应用 | 适用于: ICC/IF, WB, IHC-P |
| 种属反应性 | 与反应: Human |
| 免疫原 | Synthetic peptide derived from human Cytokeratin 17. |
| 常规说明 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C. |
| 存储溶液 | <p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituents: 50% Glycerol, 0.87% Sodium chloride, PBS</p> |
| 纯度 | Without Mg+2 and Ca+2 |
| 纯化说明 | Immunogen affinity purified |
| 克隆 | ab53707 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| 同种型 | 多克隆 |
| | IgG |

应用

The Abpromise guarantee


Abpromise™承诺保证使用ab53707于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|--------|-----------|--|
| ICC/IF | ★★★★★ (1) | Use a concentration of 1 µg/ml. |
| WB | | 1/500 - 1/1000. Detects a band of approximately 48 kDa (predicted molecular weight: 48 kDa). |
| IHC-P | | 1/50 - 1/100. |

| | |
|-------|--|
| 靶标 | |
| 功能 | May play a role in the formation and maintenance of various skin appendages, specifically in determining shape and orientation of hair. May be a marker of basal cell differentiation in complex epithelia and therefore indicative of a certain type of epithelial "stem cells". May act as an autoantigen in the immunopathogenesis of psoriasis, with certain peptide regions being a major target for autoreactive T-cells and hence causing their proliferation. Required for the correct growth of hair follicles, in particular for the persistence of the anagen (growth) state. Modulates the function of TNF-alpha in the specific context of hair cycling. Regulates protein synthesis and epithelial cell growth through binding to the adapter protein SFN and by stimulating Akt/mTOR pathway. Involved in tissue repair. |
| 组织特异性 | Expressed in the outer root sheath and medulla region of hair follicle specifically from eyebrow and beard, digital pulp, nail matrix and nail bed epithelium, mucosal stratified squamous epithelia and in basal cells of oral epithelium, palmoplantar epidermis and sweat and mammary glands. Also expressed in myoepithelium of prostate, basal layer of urinary bladder, cambial cells of sebaceous gland and in exocervix (at protein level). |
| 疾病相关 | Defects in KRT17 are a cause of pachyonychia congenita type 2 (PC2) [MIM:167210]; also known as pachyonychia congenita Jackson-Lawler type. PC2 is an autosomal dominant ectodermal dysplasia characterized by hypertrophic nail dystrophy resulting in onchyogryposis (thickening and increase in curvature of the nail), palmoplantar keratoderma and hyperhidrosis, follicular hyperkeratosis, multiple epidermal cysts, absent/sparse eyebrow and body hair, and by the presence of natal teeth. Defects in KRT17 are a cause of steatocystoma multiplex (SM) [MIM:184500]. SM is a disease characterized by round or oval cystic tumors widely distributed on the back, anterior trunk, arms, scrotum, and thighs. Note=KRT16 and KRT17 are coexpressed only in pathological situations such as metaplasias and carcinomas of the uterine cervix and in psoriasis vulgaris. |
| 序列相似性 | Belongs to the intermediate filament family. |
| 细胞定位 | Cytoplasm. |


图片



Peptide - +

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Cytokeratin 17 antibody - Cytoskeleton Marker (ab53707)

ab53707 at 1/50 dilution staining Cytokeratin 17 in human breast carcinoma by Immunohistochemistry, Paraffin embedded tissue, in the absence or presence of the immunising peptide.



Western blot - Anti-Cytokeratin 17 antibody - Cytoskeleton Marker (ab53707)


All lanes : Anti-Cytokeratin 17 antibody - Cytoskeleton Marker (ab53707) at 1/500 dilution

Lane 1 : HuvEc cell extract

Lane 2 : HuvEc cell extract with immunising peptide

Predicted band size: 48 kDa

Observed band size: 48 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Cytokeratin 17 antibody - Cytoskeleton Marker (ab53707)

ICC/IF image of ab53707 stained HeLa cells. The cells were 4% formaldehyde fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab53707, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors