


Anti-Dab1 antibody ab111684

[1 Abreviews](#) [4 References](#) [4 图像](#)

概述

产品名称	Anti-Dab1抗体
描述	兔多克隆抗体to Dab1
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Chicken, Pig, Zebrafish 
免疫原	Recombinant fragment corresponding to Human Dab1 aa 1-241. Recombinant fragment corresponding to a region within amino acids 1 and 241 of DAB1 Database link: O75553
阳性对照	WB: Mouse and rat brain cell lysate. IHC-P: Mouse duodenum tissue. ICC/IF: HepG2 cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.00 Preservative: 0.025% Proclin 300 Constituents: 79% PBS, 20% Glycerol (glycerin, glycerine)
纯度	Immunogen affinity purified
纯化说明	ab111684 is purified by antigen-affinity chromatography.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab111684于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/500 - 1/3000. Predicted molecular weight: 60 kDa.
IHC-P		1/100 - 1/1000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		1/100 - 1/1000.

靶标


功能	Adapter molecule functioning in neural development. May regulate SIAH1 activity.
序列相似性	Contains 1 PID domain.
结构域	The PID domain specifically binds to the Asn-Pro-Xaa-Tyr(P) motif found in many tyrosine-phosphorylated proteins.
翻译后修饰	Phosphorylated on Tyr-198 and Tyr-220 upon reelin induction in embryonic neurons (By similarity). Also phosphorylated on Ser-524 independently of reelin signaling.

图片



Immunocytochemistry/ Immunofluorescence - Anti-Dab1 antibody (ab111684)

HepG2 cells stained for DAB1 (green) using ab111684 at 1/500 dilution in ICC/IF.



Western blot - Anti-Dab1 antibody (ab111684)

All lanes : Anti-Dab1 antibody (ab111684) at 1/1000 dilution


Lane 1 : Mouse brain tissue extract

Lane 2 : Rat brain tissue extract

Lysates/proteins at 50 µg per lane.


Predicted band size: 60 kDa

7.5% SDS-PAGE



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Dab1 antibody (ab111684)

Paraffin embedded mouse duodenum tissue stained for DAB1 using ab111684 at 1/500 dilution in immunohistochemical analysis.



Anti-Dab1 antibody (ab111684) at 1/3000 dilution + Mouse brain cell lysate at 50 µg

Predicted band size: 60 kDa

7.5% SDS-PAGE.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors