

Anti-DAD1 antibody ab23836

★★★★★ [1 Abreviews](#) [2 References](#) [2 图像](#)

概述

产品名称	Anti-DAD1抗体
描述	兔多克隆抗体to DAD1
宿主	Rabbit
经测试应用	适用于: WB, ICC
种属反应性	与反应: Human
免疫原	Synthetic peptide corresponding to a 14 amino acid sequence within the C terminal region of DAD 1 (Human).
阳性对照	HepG2 cell lysates.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium azide Constituent: PBS
纯度	Protein A purified
纯化说明	Purified by affinity chromatography.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab23836于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (1)	Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 22 kDa (predicted molecular weight: 12 kDa).
ICC		Use at an assay dependent dilution.

靶标

功能	Component of the N-oligosaccharyl transferase enzyme which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). Loss of the DAD1 protein triggers apoptosis.
序列相似性	Belongs to the DAD/OST2 family.
细胞定位	Membrane.


图片



Lane 1 : Anti-DAD1 antibody (ab23836) at 0.5 µg/ml
Lane 2 : Anti-DAD1 antibody (ab23836) at 1 µg/ml
Lane 3 : Anti-DAD1 antibody (ab23836) at 2 µg/ml

All lanes : HepG2 whole cell lysate

Predicted band size: 12 kDa
Observed band size: 22 kDa



ab23836 staining DAD1 in HepG2 cells by Immunocytochemistry.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors